

5.4.4.2.	CELL-BASED ASSAYS	63
5.5.	SCREENING ASSAYS FOR COMPOUNDS THAT INTERACT WITH THE TARGET GENE PRODUCT AND/OR MODULATE TARGET GENE EXPRESSION	66
5.5.1.	IN VITRO SCREENING ASSAYS FOR COMPOUNDS THAT BIND TO THE TARGET GENE PRODUCT	67
5.5.2.	ASSAYS FOR CELLULAR OR EXTRACELLULAR PROTEINS THAT INTERACT WITH THE TARGET GENE PRODUCT	70
5.5.3.	ASSAYS FOR COMPOUNDS THAT INTERFERE WITH INTERACTION BETWEEN TARGET GENE PRODUCT AND OTHER COMPOUNDS	70
5.5.4.	ASSAYS FOR AMELIORATION OF CARDIOVASCULAR DISEASE SYMPTOMS	80
5.5.5.	MONITORING OF EFFECTS DURING CLINICAL TRIALS	82
5.5.6.	ASSAYS FOR COMPOUNDS THAT MODULATE EXPRESSION OF TARGET GENES	84
5.6.	COMPOUNDS AND METHODS FOR TREATMENT OF CARDIOVASCULAR DISEASE	84
5.6.1.	COMPOUNDS THAT INHIBIT EXPRESSION, SYNTHESIS OR ACTIVITY OF MUTANT TARGET GENE ACTIVITY	86
5.6.1.1.	87
5.6.1.2.	ANTIBODIES FOR TARGET GENE PRODUCTS	96
5.6.2.	METHODS FOR RESTORING OR ENHANCING TARGET GENE ACTIVITY	97
5.7.	PHARMACEUTICAL PREPARATIONS AND METHODS OF ADMINISTRATION	99
5.7.1.	EFFECTIVE DOSE	99
5.7.2.	FORMULATIONS AND USE	100
5.8.	DIAGNOSIS OF CARDIOVASCULAR DISEASE ABNORMALITIES	103
5.8.1.	DETECTION OF FINGERPRINT GENE NUCLEIC ACIDS	103
5.8.2.	DETECTION OF FINGERPRINT GENE PEPTIDES	106
5.8.3.	IMAGING CARDIOVASCULAR DISEASE CONDITIONS	111
6.	EXAMPLE: IDENTIFICATION OF GENES DIFFERENTIALLY	

	Page
EXPRESSED IN RESPONSE TO PARADIGM A: IN VITRO FOAM CELL PARADIGM	113
6.1. MATERIALS AND METHODS	114
6.1.1. CELL ISOLATION AND CULTURING	114
6.1.2. ANALYSIS OF PARADIGM MATERIAL	116
6.1.3. CHROMOSOMAL LOCALIZATION OF TARGET GENES	122
7. EXAMPLE: IDENTIFICATION OF GENES DIFFERENTIALLY EXPRESSED IN RESPONSE TO PARADIGM B: IN VIVO MONOCYTES	123
7.1. MATERIALS AND METHODS	125
7.1.1. IN VIVO CHOLESTEROL STUDIES	125
7.1.2. PRELIMINARY DETECTION SYSTEM	127
7.1.3. TRANSGENIC ApoE-DEFICIENT MOUSE EXPRESSING HUMAN bcl-2	128
7.2. RESULTS	129
8. EXAMPLE: IDENTIFICATION OF GENES DIFFERENTIALLY EXPRESSED IN RESPONSE TO PARADIGM C: IL-1 INDUCTION OF ENDOTHELIAL CELLS	131
8.1. MATERIALS AND METHODS	132
8.2. RESULTS	133
9. EXAMPLE: IDENTIFICATION OF GENES DIFFERENTIALLY EXPRESSED IN RESPONSE TO PARADIGM D: ENDOTHELIAL CELL SHEAR STRESS	134
9.1. MATERIALS AND METHODS	137
9.2. RESULTS	139
10. EXAMPLE: USE OF GENES UNDER PARADIGM A AS SURROGATE MARKERS IN CLINICAL TRIALS	143
10.1. TREATMENT OF PATIENTS AND CELL ISOLATION	144
10.2. ANALYSIS OF SAMPLES	144
11. EXAMPLE: IMAGING OF A CARDIOVASCULAR DISEASE CONDITION	144
11.1. MONOCLONAL CONJUGATED ANTIBODIES	145
11.2. ADMINISTRATION AND DETECTION OF IMAGING	

	Page
AGENTS	145
12. EXAMPLE: SCREENING FOR LIGANDS OF THE rchd523 GENE PRODUCT AND ANTAGONISTS OF rchd523 GENE PRODUCT-LIGAND INTERACTION	145
13. POLYCLONAL ANTIBODIES TO TARGET GENE PEPTIDE SEQUENCES	146
14. LOCALIZATION OF NOVEL GENES BY <i>IN SITU</i> HYBRIDIZATION	148
14.1 Methods	149
14.2 Results	150
15. EXAMPLE: THE RCHD534 AND FCHD540 GENE PRODUCTS INTERACT	150
15.1 MATERIALS AND METHODS	152
1. YEAST STRAINS, MEDIA, AND MICROBIOLOGICAL TECHNIQUES	152
2. PLASMID AND YEAST STRAIN CONSTRUCTION	152
15.1.4 PAPER FILTER BETA-GALACTOSIDASE ASSAYS	153
15.2 RESULTS	153
15.2.1 STRONG PHYSICAL INTERACTION OF RCHD534 AND FCHD540 MEASURED BY TWO- HYBRID ASSAY	153
15.2.2 IDENTIFICATION OF PROTEINS THAT PHYSICALLY INTERACT WITH FCHD540	154
15.2.3 RETRANSFORMATION AND SPECIFICITY TESTING OF TCHV03A AND TCHVR4A	155
15.3 FURTHER ANALYSIS OF RCHD534 AND FCHD540 FUNCTION	157
15.3.1 CHROMOSOMAL LOCALIZATION	157
15.3.2 TISSUE EXPRESSION PATTERNS	158
15.3.3. CELLULAR LOCALIZATION	159
15.3.4. PROTEIN INTERACTIONS IN HUMAN CELLS	159
15.3.5 EFFECT OF EXPRESSION ON TGF-B SIGNALLING	160
16. EXAMPLE: THE RCHD534-LONG PROTEIN	162

0640E0" 9824E060

	<u>Page</u>
16.1 IDENTIFICATION AND CHARACTERIZATION OF THE RCHD534-LONG SPLICEFORM AND PROTEIN . . .	162
16.2 TGF- β SIGNALLING INHIBITORY ACTIVITY OF THE RCHD534-LONG PROTEIN	163
17. EXAMPLE: ANTISENSE AND RIBOZYME MOLECULES FOR INHIBITION OF RCHD534 AND FCHD540 EXPRESSION	165
17.1. ANTISENSE MOLECULES	165
17.2. RIBOZYME MOLECULES	167
18. DEPOSIT OF MICROORGANISMS	169

COMPOSITIONS AND METHODS FOR THE TREATMENT
AND DIAGNOSIS OF CARDIOVASCULAR DISEASE

5 This application is a continuation-in-part of co-
pending application serial number 08/870,434, filed June 6,
1997, which is a continuation-in-part of co-pending
application serial number 08/799,910, filed February 13,
1997, which claims the benefit under 35 U.S.C. § 119(e) of
provisional application serial number 60/011,787, filed
10 February 16, 1996, now abandoned; and this application is a
continuation-in-part of co-pending application serial number
08/599,654, filed February 9, 1996, which is a continuation-
in-part of co-pending application serial number 08/485,573,
filed June 7, 1995, which is a continuation-in-part of co-
15 pending application serial number 08/386,844, filed February
10, 1995, each of which is hereby incorporated by reference
in its entirety.

1. INTRODUCTION

20 The present invention relates to methods and
compositions for the treatment and diagnosis of
cardiovascular disease, including, but not limited to,
atherosclerosis, ischemia/reperfusion, hypertension,
restenosis, and arterial inflammation. The present invention
25 further relates to screening methods to identify compositions
and their therapeutic use for the treatment of fibro-
proliferative and oncogenic disorders, including diabetic
retinopathy, arteriosclerosis, angiogenesis, inflammation,
fibrosis, tumor growth and vascularization. Genes which are
30 differentially expressed in cardiovascular or oncogenic
disease states, relative to their expression in normal, or
non-disease states are identified. Genes are also identified
via the ability of their gene products to interact with other
gene products involved in cardiovascular or oncogenic
35 disease. The genes identified may be used diagnostically or
as targets for therapeutic intervention. In this regard, the
present invention provides methods for the identification and

pathway by which native LDL (nLDL) is taken up by nLDL specific receptors, the scavenger pathway of uptake is not regulated by the monocytes.

These lipid-filled monocytes are called foam cells,
5 and are the major constituent of the fatty streak.

Interactions between foam cells and the endothelial and SMCs which surround them lead to a state of chronic local inflammation which can eventually lead to smooth muscle cell proliferation and migration, and the formation of a fibrous
10 plaque. Such plaques occlude the blood vessel concerned and thus restrict the flow of blood, resulting in ischemia.

Ischemia is a condition characterized by a lack of oxygen supply in tissues of organs due to inadequate perfusion. Such inadequate perfusion can have number of
15 natural causes, including atherosclerotic or restenotic lesions, anemia, or stroke, to name a few. Many medical interventions, such as the interruption of the flow of blood during bypass surgery, for example, also lead to ischemia. In addition to sometimes being caused by diseased
20 cardiovascular tissue, ischemia may sometimes affect cardiovascular tissue, such as in ischemic heart disease. Ischemia may occur in any organ, however, that is suffering a lack of oxygen supply.

The most common cause of ischemia in the heart is
25 atherosclerotic disease of epicardial coronary arteries. By reducing the lumen of these vessels, atherosclerosis causes an absolute decrease in myocardial perfusion in the basal state or limits appropriate increases in perfusion when the demand for flow is augmented. Coronary blood flow can also
30 be limited by arterial thrombi, spasm, and, rarely, coronary emboli, as well as by ostial narrowing due to luetic aortitis. Congenital abnormalities, such as anomalous origin of the left anterior descending coronary artery from the pulmonary artery, may cause myocardial ischemia and
35 infarction in infancy, but this cause is very rare in adults. Myocardial ischemia can also occur if myocardial oxygen demands are abnormally increased, as in severe ventricular

hypertrophy due to hypertension or aortic stenosis. The latter can be present with angina that is indistinguishable from that caused by coronary atherosclerosis. A reduction in the oxygen-carrying capacity of the blood, as in extremely
5 severe anemia or in the presence of carboxy-hemoglobin, is a rare cause of myocardial ischemia. Not infrequently, two or more causes of ischemia will coexist, such as an increase in oxygen demand due to left ventricular hypertrophy and a reduction in oxygen supply secondary to coronary
10 atherosclerosis.

The principal surgical approaches to the treatment of ischemic atherosclerosis are bypass grafting, endarterectomy, and percutaneous transluminal angioplasty (PCTA). The failure rate after these approaches due to
15 restenosis, in which the occlusions recur and often become even worse, is extraordinarily high (30-50%). It appears that much of the restenosis is due to further inflammation, smooth muscle accumulation, and thrombosis.

Very recently, a modified balloon angioplasty
20 approach was used to treat arterial restenosis in pigs by gene therapy (Ohno et al., 1994, Science 265: 781-784). A specialized catheter was used to introduce a recombinant adenovirus carrying the gene encoding thymidine kinase (tk) into the cells at the site of arterial blockage.
25 Subsequently, the pigs were treated with ganciclovir, a nucleoside analog which is converted by tk into a toxic form which kills cells when incorporated into DNA. Treated animals had a 50% to 90% reduction in arterial wall thickening without any observed local or systemic toxicities.

30 Because of the presumed role of the excessive inflammatory-fibroproliferative response in atherosclerosis and ischemia, a number of researchers have investigated, in the context of arterial injury, the expression of certain factors involved in inflammation, cell recruitment and
35 proliferation. These factors include growth factors, cytokines, and other chemicals, including lipids involved in

Recently, cytomegalovirus (CMV) has been implicated in restenosis as well as atherosclerosis in general (Speir, et al., 1994, Science 265: 391-394). It was observed that the CMV protein IE84 apparently predisposes smooth muscle
5 cells to increased growth at the site of restenosis by combining with and inactivating p53 protein, which is known to suppress tumors in its active form.

The foregoing studies are aimed at defining the role of particular gene products presumed to be involved in
10 the excessive inflammatory-fibroproliferative response leading to atherosclerotic plaque formation. However, such approaches cannot identify the full panoply of gene products that are involved in the disease process, much less identifying those which may serve as therapeutic targets for
15 the diagnosis and treatment of various forms of cardiovascular disease.

3. SUMMARY OF THE INVENTION

The present invention relates to methods and
20 compositions for the treatment and diagnosis of cardiovascular disease, including but not limited to, atherosclerosis, ischemia/reperfusion, hypertension, restenosis, and arterial inflammation. Specifically, genes are identified and described which are differentially
25 expressed in cardiovascular disease states, relative to their expression in normal, or non-cardiovascular disease states.

The present invention further relates to screening methods to identify compositions and their therapeutic use for the treatment of fibroproliferative and oncogenic
30 disorders, including diabetic retinopathy, cancer, tumorigenesis, vascularization of tumors, angiogenesis atherosclerosis inflammation and fibrosis.

"Differential expression", as used herein, refers to both quantitative as well as qualitative differences in
35 the genes' temporal and/or tissue expression patterns. Differentially expressed genes may represent "fingerprint genes," and/or "target genes." "Fingerprint gene," as used

herein, refers to a differentially expressed gene whose expression pattern may be utilized as part of a prognostic or diagnostic cardiovascular disease evaluation, or which, alternatively, may be used in methods for identifying

5 compounds useful for the treatment of cardiovascular disease.

"Target gene", as used herein, refers to a differentially expressed gene involved in cardiovascular disease such that modulation of the level of target gene expression or of target gene product activity may act to ameliorate a

10 cardiovascular disease condition. Compounds that modulate target gene expression or activity of the target gene product can be used in the treatment of cardiovascular disease.

Further, "pathway genes" are defined via the ability of their products to interact with other gene

15 products involved in cardiovascular disease. Pathway genes may also exhibit target gene and/or fingerprint gene characteristics. Although the genes described herein may be differentially expressed with respect to cardiovascular disease, and/or their products may interact with gene

20 products important to cardiovascular disease, the genes may also be involved in mechanisms important to additional cardiovascular processes.

The invention includes the products of such fingerprint, target, and pathway genes, as well as antibodies
25 to such gene products. Furthermore, the engineering and use of cell- and animal-based models of cardiovascular disease to which such gene products may contribute are also described.

The present invention encompasses methods for prognostic and diagnostic evaluation of cardiovascular
30 disease conditions, and for the identification of subjects exhibiting a predisposition to such conditions. Furthermore, the invention provides methods for evaluating the efficacy of drugs, and monitoring the progress of patients, involved in clinical trials for the treatment of cardiovascular disease.

35 The invention also provides methods for the identification of compounds that modulate the expression of genes or the activity of gene products involved in

cardiovascular disease, as well as methods for the treatment of cardiovascular disease which may involve the administration of such compounds to individuals exhibiting cardiovascular disease symptoms or tendencies.

5 The invention also provides methods for the identification of compounds that modulate the expression of genes or the activity of gene products involved in fibroproliferative or oncogenic disorders, including tumorigenesis and the vascularization of tumors.

10 In addition, the invention encompasses methods for treating cardiovascular disease and fibroproliferative or oncogenic disorders by administering compounds and other substances that modulate the overall activity of the target gene products. Compounds and other substances can effect
15 such modulation either on the level of target gene expression or target protein activity.

 The invention is based, in part, on systematic search strategies involving in vivo and in vitro cardiovascular disease paradigms coupled with sensitive and
20 high throughput gene expression assays. In contrast to approaches that merely evaluate the expression of a given gene product presumed to play a role in a disease process, the search strategies and assays used herein permit the identification of all genes, whether known or novel, that are
25 expressed or repressed in the disease condition, as well as the evaluation of their temporal regulation and function during disease progression. This comprehensive approach and evaluation permits the discovery of novel genes and gene products, as well as the identification of an array of genes
30 and gene products (whether novel or known) involved in novel pathways that play a major role in the disease pathology. Thus, the invention allows one to define targets useful for diagnosis, monitoring, rational drug screening and design, and/or other therapeutic intervention.

35 In the working examples described herein, eight novel human genes are identified that are demonstrated to be differentially expressed in different cardiovascular disease

states. Additionally, the differential expression of four previously identified human genes is described. The identification of these genes and the characterization of their expression in particular disease states provide newly
5 identified roles in cardiovascular disease for both the novel genes and the known genes.

Bcl-2 and glutathione peroxidase are the products of known genes that are shown herein to be down regulated in monocytes of patients exposed to an atherogenic high fat/high
10 cholesterol diet. Furthermore, counteracting the down-regulation of bcl-2 under atherogenic conditions, as described herein, may ameliorate atherosclerosis. Accordingly, methods are provided for the diagnosis, monitoring in clinical trials, and treatment of
15 cardiovascular disease based upon the discoveries herein regarding the expression patterns of bcl-2 and glutathione peroxidase. Because these two genes were known to be involved in preventing apoptosis, the discovery of their down-regulation under atherogenic conditions provides a
20 novel, positive correlation between apoptosis and atherogenesis. Accordingly, methods provided herein for diagnosing, monitoring, and treating cardiovascular disease may also be based on a number of genes involved in the apoptotic pathway, including but not limited to ICE (IL-1
25 converting enzyme); Bad; BAG-1 (Bcl-2 associated athanogene 1, Takayama et al., 1995, Cell 80: 279-284); BAX (Bcl-2 associated X protein, Oltvai et al., 1993, Cell 74: 609-619); BclX_L (Boise, et al., 1993, Cell 74: 597-608); BAK (Bcl-2 antagonist killer, Farrow et al., 1995. Nature 374: 631-733);
30 and Bcl-X_s (Tsujimoto et al., 1984, Science 226: 1097-1099). The cardiovascular diseases that may be so diagnosed, monitored in clinical trials, and treated include but are not limited to atherosclerosis, ischemia/reperfusion, and restenosis.

35 rchd005, rchd024, rchd032, and rchd036 are newly identified genes that are each up-regulated in endothelial cells treated with IL-1. Accordingly, methods are provided

for the diagnosis, monitoring in clinical trials, and treatment of cardiovascular disease based upon the discoveries herein regarding the expression patterns of rchd005, rchd024, rchd032, and rchd036.

- 5 Cyclooxygenase II (COX II), also known as endoperoxide synthase, and Manganese Superoxide Dismutase (MnSOD) are known genes, and rchd502, rchd523, rchd528, and rchd534 are newly identified genes, that are each up-regulated in endothelial cells subjected to shear stress.
- 10 Accordingly, methods are provided for the diagnosis, monitoring in clinical trials, screening for therapeutically effective compounds, and treatment of cardiovascular disease based upon the discoveries herein regarding the expression patterns of COX II, MnSOD, rchd502, rchd523, rchd528, and
- 15 rchd534.

- More specifically, each of these genes is up-regulated either by IL-1 (rchd005, rchd024, rchd032, and rchd036) or by shear stress (COX II, MnSOD, rchd502, rchd523, rchd528, and rchd534). For those genes that have a causative
- 20 effect on the disease conditions treatment methods can be designed to reduce or eliminate their expression, particularly in endothelial cells. Alternatively, treatment methods include inhibiting the activity of the protein products of these genes. For those genes that have a
- 25 protective effect in responding to disease conditions, treatment methods can be designed for enhancing the activity of the products of such genes.

- In either situation, detecting expression of these genes in excess of normal expression provides for the
- 30 diagnosis of cardiovascular disease. Furthermore, in testing the efficacy of compounds during clinical trials, a decrease in the level of the expression of these genes corresponds to a return from a disease condition to a normal state, and thereby indicates a positive effect of the compound. The
- 35 cardiovascular diseases that may be so diagnosed, monitored in clinical trials, and treated include but are not limited

to atherosclerosis, ischemia/reperfusion, hypertension, restenosis, and arterial inflammation.

Membrane bound target gene products containing extracellular domains can be a particularly useful target for treatment methods as well as diagnostic and clinical monitoring methods. The rchd523 gene, for example, encodes a transmembrane protein, which contains seven transmembrane domains and, therefore, can be readily contacted by other compounds on the cell surface. Accordingly, natural ligands, derivatives of natural ligands, and antibodies that bind to the rchd523 gene product can be utilized to inhibit its activity, or alternatively, to target the specific destruction of cells that are in the disease state. Furthermore, the extracellular domains of the rchd523 gene product provide especially efficient screening systems for identifying compounds that bind to the rchd523 gene product. Compounds that bind the receptor domain of the rchd523 gene product, for example, can be identified by their ability to mobilize Ca^{2+} and thereby produce a fluorescent signal, as described in Section 5.5.1, below.

Such an assay system can also be used to screen and identify antagonists of the interaction between the rchd523 gene product and ligands that bind to the rchd523 gene product. For example, the compounds can compete with the endogenous (i.e., natural) ligand for the rchd523 gene product. The resulting reduction in the amount of ligand-bound rchd523 gene transmembrane protein will modulate the activity of disease state cells, such as endothelial cells. Soluble proteins or peptides, such as peptides comprising one or more of the extracellular domains, or portions and/or analogs thereof of the rchd523 gene product, including, for example, soluble fusion proteins such as Ig-tailed fusion proteins, can be particularly useful for this purpose.

Similarly, antibodies that are specific to one or more of the extracellular domains of the rchd523 product provide for the ready detection of this target gene product in diagnostic tests or in clinical test monitoring.

Accordingly, endothelial cells can be treated, either in vivo or in vitro, with such a labeled antibody to determine the disease state of endothelial cells. Because the rchd523 gene product is up-regulated in endothelial cells under shear
5 stress, its detection positively corresponds with cardiovascular disease.

Such methods for treatment, diagnosis, and clinical test monitoring which use the rchd523 gene product as described above can also be applied to other target genes
10 that encode transmembrane gene products, including but not limited to rchd502, which each contains 12 transmembrane domains, and rchd528, which contains one transmembrane domain in addition to its extracellular domain.

The invention is based in part on the
15 identification of novel protein-protein interactions of the rchd534 protein with itself and with the fchd540 protein, encoded by the fchd540 gene, as well as interactions of the rchd534 protein or the fchd540 protein with other protein members of the TGF- β signalling pathway. The fchd540 gene
20 and protein were described in co-pending Application No. 08/799,910, filed February 13, 1997, which is hereby incorporated by reference in its entirety. Screening methods are provided for identifying compounds and other substances for treating cardiovascular disease by assaying their ability
25 to inhibit these interactions. Furthermore, methods are provided for identifying compounds and other substances that enhance the TGF- β response by modulating the expression of the rchd534 or fchd540 genes or the activity of their gene products. In addition, methods are provided for treating
30 cardiovascular disease by administering compounds and other substances that inhibit these protein interactions.

In addition, the invention is based in part on the identification of the endothelial cell specific expression pattern of the rchd534 and fchd540 genes, whose protein
35 products inhibit the TGF- β response. The fchd540 gene has been mapped to regions of the human genome that have been implicated in the pathogenesis of several human malignancies.

The invention is further based on the finding that these genes and mutants thereof may be used to modulate TGF- β induced signalling in endothelial cells. Accordingly, the rchd534 and fchd540 genes may be targets for intervention in
5 a variety of inflammatory and fibroproliferative disorders that involve endothelial cells, including, but not limited to, oncology related disorders, disorders related to vascularization, such as cancer angiogenesis, inflammation, and fibrosis.

- 10 Both fchd540 and rchd534 are up-regulated in response to laminar shear stress and are specifically expressed in vascular tissue. These findings combined with the observations that both the fchd540 and rchd534 proteins specifically inhibit TGF- β signalling and that the fchd540
15 and rchd534 genes are located in an area of the human genome implicated in the pathogenesis of several human malignancies indicates that they are excellent and specific targets for therapeutic intervention in the treatment of fibroproliferative and oncogenic disorders including
20 tumorigenesis and vascularization.

- The invention is also based in part on the discovery that the rchd534 gene encodes a second protein, in addition to the rchd534 protein described in co-pending application number 08/485,573. Specifically, the rchd534
25 gene also encodes the rchd534-long protein. The rchd534 protein and the rchd534-long protein are encoded by alternative spliceoforms of rchd534 mRNA. Thus, the rchd534 protein is encoded by the short rchd534 spliceoform, and the rchd534-long protein is encoded by the long rchd534
30 spliceoform. The rchd534 and rchd534-long protein each has a common C-terminal domain (MH2 domain). The rchd534-long protein contains an N-terminal MH1 domain that is not present in the rchd534 protein. Like rchd534 and fchd540, the rchd534-long protein inhibits TGF- β signalling.

- 35 The examples presented in Sections 6-9, below, demonstrate the use of the cardiovascular disease paradigms

of the invention to identify cardiovascular disease target genes.

The example presented in Section 10, below, demonstrates the use of fingerprint genes in diagnostics and as surrogate markers for testing the efficacy of candidate drugs in basic research and in clinical trials.

The example presented in Section 11, below, demonstrates the use of fingerprint genes, particularly rchd523, in the imaging of a diseased cardiovascular tissue.

The example presented in Section 12, below, demonstrates the use of target genes, particularly rchd523, in screening for ligands of target gene product receptor domains, as well as antagonists of the ligand-receptor interaction.

The example presented in Section 15, below, demonstrates the interaction of two target gene products, the rchd534 and fchd540 proteins, and the further characterization of their roles in oncology, angiogenesis, cardiovascular disease and the TGF- β signalling pathway.

The example presented in Section 16, below, describes the discovery of the novel rchd534-long protein and demonstrates that the rchd534-long protein inhibits TGF- β signalling, as do the rchd534 and fchd540 proteins.

4. DESCRIPTION OF THE FIGURES

FIG.1. Band rchd005 DNA sequence. ^(SEQ ID NO: 52) The sequence was determined by sequencing the insert of pRCHD005, resulting from the ligation of amplified rchd005 sequences into the TA cloning vector.

FIG.2. Band rchd024 DNA sequence. ^(SEQ ID NO: 53) The sequence was determined by sequencing the insert of pRCHD024, resulting from the ligation of amplified rchd024 sequences into the TA cloning vector.

FIG.3. Band rchd032 DNA sequence. ^(SEQ ID NO: 54) The sequence was determined by sequencing the insert of pRCHD032, resulting from the ligation of amplified rchd032 sequences into the TA cloning vector.

FIG.4. Band rchd036 DNA sequence. ^(SEQ ID NO:55) The sequence was determined by sequencing the insert of pRCHD036, resulting from the ligation of amplified rchd036 sequences into the TA cloning vector.

FIG.5. DNA ^(SEQ ID NO:57) and encoded amino acid ^(SEQ ID NO:56) sequence of the rchd502 gene.

FIG.6. DNA ^(SEQ ID NO:58) and encoded amino acid ^(SEQ ID NO:59) sequence of the rchd523 gene.

FIG.7. DNA ^(SEQ ID NO:61) and encoded amino acid ^(SEQ ID NO:60) sequence of the rchd528 gene.

FIG.8. DNA ^(SEQ ID NO:62) and encoded amino acid ^(SEQ ID NO:63) sequence of the rchd534 cDNA encoding the rchd534 protein.

FIG. 9. DNA ^(SEQ ID NO:64) and encoded amino acid ^(SEQ ID NO:65) sequence of the rchd534-long cDNA encoding the rchd534-long protein.

FIG.10. DNA ^(SEQ ID NO:66) and encoded amino acid ^(SEQ ID NO:67) sequence of the fchd540 gene.

FIG.11. Schematic comparison of the rchd534 protein and the rchd534-long protein.

FIG.12. Northern blot analysis of expression of rchd534 mRNA (short spliceoform) under shear stress. RNA was prepared from HUVEC's that were untreated (control) and treated with shear stress for 1 hr. and 6 hr. The blot was probed with labeled rchd534 DNA.

25

5. DETAILED DESCRIPTION OF THE INVENTION

Methods and compositions for the diagnosis and treatment of cardiovascular disease, including but not limited to atherosclerosis, ischemia/reperfusion, hypertension, restenosis, and arterial inflammation, are described. Methods and compositions for the treatment of oncogenic related disorders, including tumorigenesis and the vascularization of tumors, are also described. The invention is based, in part, on the evaluation of the expression and role of all genes that are differentially expressed in paradigms that are physiologically relevant to the disease condition. This permits the definition of disease pathways

and the identification of targets in the pathway that are useful both diagnostically and therapeutically.

Genes, termed "target genes" and/or "fingerprint genes" which are differentially expressed in cardiovascular disease conditions, relative to their expression in normal, or non-cardiovascular disease conditions, are described in Section 5.4. Additionally, genes, termed "pathway genes" whose gene products exhibit an ability to interact with gene products involved in cardiovascular disease are also described in Section 5.4. Pathway genes may additionally have fingerprint and/or target gene characteristics. Methods for the identification of such fingerprint, target, and pathway genes are described in Sections 5.1, 5.2, and 5.3.

Further, the gene products of such fingerprint, target, and pathway genes are described in Section 5.4.2, antibodies to such gene products are described in Section 5.4.3, as are cell- and animal-based models of cardiovascular disease and fibroproliferative and oncogenic related disorders to which such gene products may contribute, in Section 5.4.4.

Methods for the identification of compounds which modulate the expression of genes or the activity of gene products involved in cardiovascular disease and fibroproliferative and oncogenic related disorders are described in Section 5.5. Methods for monitoring the efficacy of compounds during clinical trials are described in Section 5.5.4. Additionally described below, in Section 5.6, are methods for the treatment of cardiovascular disease.

Also discussed below, in Section 5.8, are methods for prognostic and diagnostic evaluation of cardiovascular disease and fibroproliferative and oncogenic related disorders, including the identification of subjects exhibiting a predisposition to this disease, and the imaging of cardiovascular disease conditions.

35

or decreased, in normal versus cardiovascular disease states, or under control versus experimental conditions. The degree to which expression differs in normal versus cardiovascular disease or control versus experimental states need only be
5 large enough to be visualized via standard characterization techniques, such as, for example, the differential display technique described below. Other such standard characterization techniques by which expression differences may be visualized include but are not limited to quantitative
10 RT-PCR and Northern analyses.

Differentially expressed genes may be further described as target genes and/or fingerprint genes.

"Fingerprint gene," as used herein, refers to a differentially expressed gene whose expression pattern may be
15 utilized as part of a prognostic or diagnostic cardiovascular disease evaluation, or which, alternatively, may be used in methods for identifying compounds useful for the treatment of cardiovascular disease. A fingerprint gene may also have the characteristics of a target gene.

20 "Target gene", as used herein, refers to a differentially expressed gene involved in cardiovascular disease in a manner by which modulation of the level of target gene expression or of target gene product activity may act to ameliorate symptoms of cardiovascular disease. A
25 target gene may also have the characteristics of a fingerprint gene.

A variety of methods may be utilized for the identification of genes which are involved in cardiovascular disease. These methods include but are not limited to the
30 experimental paradigms described, below, in Section 5.1.1. Material from the paradigms may be characterized for the presence of differentially expressed gene sequences as discussed, below, in Section 5.1.2.

35

5.1.1. PARADIGMS FOR THE IDENTIFICATION OF
DIFFERENTIALLY EXPRESSED GENES

One strategy for identifying genes that are involved in cardiovascular disease is to detect genes that are expressed differentially under conditions associated with the disease versus non-disease conditions. The sub-sections below describe a number of experimental systems, called paradigms, which may be used to detect such differentially expressed genes. In general, the paradigms include at least one experimental condition in which subjects or samples are treated in a manner associated with cardiovascular disease, in addition to at least one experimental control condition lacking such disease associated treatment. Differentially expressed genes are detected, as described herein, below, by comparing the pattern of gene expression between the experimental and control conditions.

Once a particular gene has been identified through the use of one such paradigm, its expression pattern may be further characterized by studying its expression in a different paradigm. A gene may, for example, be regulated one way in a given paradigm (e.g., up-regulation), but may be regulated differently in some other paradigm (e.g., down-regulation). Furthermore, while different genes may have similar expression patterns in one paradigm, their respective expression patterns may differ from one another under a different paradigm. Such use of multiple paradigms may be useful in distinguishing the roles and relative importance of particular genes in cardiovascular disease.

30 5.1.1.1. FOAM CELL PARADIGM - 1

Among the paradigms which may be utilized for the identification of differentially expressed genes involved in atherosclerosis, for example, are paradigms designed to analyze those genes which may be involved in foam cell formation. Such paradigms may serve to identify genes involved in the differentiation of this cell type, or their uptake of oxidized LDL.

oxidation of LDL which is then taken up by the monocytes. As described in sub-section 5.1.2. below, the pattern of gene expression can then be compared between these foam cells and untreated monocytes.

5

5.1.1.3. FOAM CELL PARADIGM - 3

Yet another system includes the third cell type, smooth muscle cell, that plays a critical role in atherogenesis (Navab et al., 1988, J. Clin. Invest., 82: 1853). In this system, a multilayer of human aortic smooth muscle cells was grown on a micropore filter covered with a gel layer of native collagen, and a monolayer of human aortic endothelial cells was grown on top of the collagen layer. Exposure of this coculture to human monocytes in the presence of chemotactic factor rFMLP resulted in monocyte attachment to the endothelial cells followed by migration across the endothelial monolayer into the collagen layer of the subendothelial space. This type of culture can also be treated with LDL to generate foam cells. The foam cells can then be harvested and their pattern of gene expression compared to that of untreated cells as explained below in sub-section 5.1.2.

5.1.1.4. IN VIVO MONOCYTE PARADIGM

An alternative embodiment of such paradigms for the study of monocytes, hereinafter referred to as Paradigm B, involves differential treatment of human subjects through the dietary control of lipid consumption. Such human subjects are held on a low fat/low cholesterol diet for three weeks, at which time blood is drawn, monocytes are isolated according to the methods routinely practiced in the art, and RNA is purified, as described below, in sub-section 5.1.2. These same patients are subsequently switched to a high fat /high cholesterol diet and monocyte RNA is purified again. The patients may also be fed a third, combination diet containing high fat/low cholesterol and monocyte RNA may be purified once again. The order in which patients receive the

diets may be varied. The RNA derived from patients maintained on two of the diets, or on all three diets, may then be compared and analyzed for differential gene expression as, explained below in sub-section 5.1.2.

5 The Example presented in Section 7, below, demonstrates the use of such an in vivo monocyte paradigm to identify genes which are expressed differentially in monocytes of patients maintained on an atherogenic diet versus their expression under a control diet. Such a
10 paradigm may also be used in conjunction with an in vitro preliminary detection system, as described in Section 7, below.

5.1.1.5. ENDOTHELIAL CELL - IL-1 PARADIGM

15 In addition to the detection of differential gene expression in monocytes, paradigms focusing on endothelial cells may be used to detect genes involved in cardiovascular disease. In one such paradigm, hereinafter referred to as Paradigm C, human umbilical vein endothelial cells (HUVEC's)
20 are grown in vitro. Experimental cultures are treated with human IL-1 β , a factor known to be involved in the inflammatory response, in order to mimic the physiologic conditions involved in the atherosclerotic state. Alternatively experimental HUVEC cultures may be treated with
25 lysophosphatidylcholine, a major phospholipid component of atherogenic lipoproteins or oxidized human LDL. Control cultures are grown in the absence of these compounds.

 After a certain period of exposure treatment, experimental and control cells are harvested and analyzed for
30 differential gene expression as described in sub-section 5.1.2, below. The Example presented in Section 8, below, demonstrates the use of such an IL-1 induced endothelial cell paradigm to identify sequences which are differentially expressed in treated versus control cells.

35

5.1.1.6. ENDOTHELIAL CELL - SHEAR STRESS
PARADIGM

In another paradigm involving endothelial cells, hereinafter referred to as Paradigm D, cultures are exposed
5 to fluid shear stress which is thought to be responsible for the prevalence of atherosclerotic lesions in areas of unusual circulatory flow. Unusual blood flow also plays a role in the harmful effects of ischemia/reperfusion, wherein an organ receiving inadequate blood supply is suddenly reperfused with
10 an overabundance of blood when the obstruction is overcome.

Cultured HUVEC monolayers are exposed to laminar shear stress by rotating the culture in a specialized apparatus containing liquid culture medium (Nagel et al., 1994, J. Clin. Invest. 94: 885-891). Static cultures grown
15 in the same medium serve as controls. After a certain period of exposure to shear stress, experimental and control cells are harvested and analyzed for differential gene expression as described in sub-section 5.1.2, below. The Example presented in Section 9, below, demonstrates the use of such a
20 shear stressed endothelial cell paradigm to identify sequences which are differentially expressed in exposed versus control cells.

In all such paradigms designed to identify genes which are involved in cardiovascular disease, including but
25 not limited to those described above in Sections 5.1.1.1 through 5.1.1.6, compounds such as drugs known to have an ameliorative effect on the disease symptoms may be incorporated into the experimental system. Such compounds may include known therapeutics, as well as compounds that are
30 not useful as therapeutics due to their harmful side effects. Test cells that are cultured as explained in the paradigms described in Sections 5.1.1.1 through 5.1.1.6, for example, may be exposed to one of these compounds and analyzed for differential gene expression with respect to untreated cells,
35 according to the methods described below in Section 5.1.2. In principle, according to the particular paradigm, any cell

differential screening (Tedder, T.F. et al., 1988, Proc. Natl. Acad. Sci. USA 85:208-212), subtractive hybridization (Hedrick, S.M. et al., 1984, Nature 308:149-153; Lee, S.W. et al., 1984, Proc. Natl. Acad. Sci. USA 88:2825), and,

5 preferably, differential display (Liang, P., and Pardee, A.B., 1993, U.S. Patent No. 5,262,311, which is incorporated herein by reference in its entirety), may be utilized to identify nucleic acid sequences derived from genes that are differentially expressed.

10 Differential screening involves the duplicate screening of a cDNA library in which one copy of the library is screened with a total cell cDNA probe corresponding to the mRNA population of one cell type while a duplicate copy of the cDNA library is screened with a total cDNA probe
15 corresponding to the mRNA population of a second cell type. For example, one cDNA probe may correspond to a total cell cDNA probe of a cell type derived from a control subject, while the second cDNA probe may correspond to a total cell cDNA probe of the same cell type derived from an experimental
20 subject. Those clones which hybridize to one probe but not to the other potentially represent clones derived from genes differentially expressed in the cell type of interest in control versus experimental subjects.

Subtractive hybridization techniques generally
25 involve the isolation of mRNA taken from two different sources, e.g., control and experimental tissue, the hybridization of the mRNA or single-stranded cDNA reverse-transcribed from the isolated mRNA, and the removal of all hybridized, and therefore double-stranded, sequences. The
30 remaining non-hybridized, single-stranded cDNAs, potentially represent clones derived from genes that are differentially expressed in the two mRNA sources. Such single-stranded cDNAs are then used as the starting material for the construction of a library comprising clones derived from
35 differentially expressed genes.

The differential display technique describes a procedure, utilizing the well known polymerase chain reaction

8640E0" 9824E050

(PCR; the experimental embodiment set forth in Mullis, K.B., 1987, U.S. Patent No. 4,683,202) which allows for the identification of sequences derived from genes which are differentially expressed. First, isolated RNA is reverse-
5 transcribed into single-stranded cDNA, utilizing standard techniques which are well known to those of skill in the art. Primers for the reverse transcriptase reaction may include, but are not limited to, oligo dT-containing primers, preferably of the reverse primer type of oligonucleotide
10 described below. Next, this technique uses pairs of PCR primers, as described below, which allow for the amplification of clones representing a random subset of the RNA transcripts present within any given cell. Utilizing different pairs of primers allows each of the mRNA
15 transcripts present in a cell to be amplified. Among such amplified transcripts may be identified those which have been produced from differentially expressed genes.

The reverse oligonucleotide primer of the primer pairs may contain an oligo dT stretch of nucleotides,
20 preferably eleven nucleotides long, at its 5' end, which hybridizes to the poly(A) tail of mRNA or to the complement of a cDNA reverse transcribed from an mRNA poly(A) tail. Second, in order to increase the specificity of the reverse primer, the primer may contain one or more, preferably two,
25 additional nucleotides at its 3' end. Because, statistically, only a subset of the mRNA derived sequences present in the sample of interest will hybridize to such primers, the additional nucleotides allow the primers to amplify only a subset of the mRNA derived sequences present
30 in the sample of interest. This is preferred in that it allows more accurate and complete visualization and characterization of each of the bands representing amplified sequences.

The forward primer may contain a nucleotide
35 sequence expected, statistically, to have the ability to hybridize to cDNA sequences derived from the tissues of interest. The nucleotide sequence may be an arbitrary one,

054020-030498

and the length of the forward oligonucleotide primer may range from about 9 to about 13 nucleotides, with about 10 nucleotides being preferred. Arbitrary primer sequences cause the lengths of the amplified partial cDNAs produced to
5 be variable, thus allowing different clones to be separated by using standard denaturing sequencing gel electrophoresis.

PCR reaction conditions should be chosen which optimize amplified product yield and specificity, and, additionally, produce amplified products of lengths which may
10 be resolved utilizing standard gel electrophoresis techniques. Such reaction conditions are well known to those of skill in the art, and important reaction parameters include, for example, length and nucleotide sequence of oligonucleotide primers as discussed above, and annealing and
15 elongation step temperatures and reaction times.

The pattern of clones resulting from the reverse transcription and amplification of the mRNA of two different cell types is displayed via sequencing gel electrophoresis and compared. Differences in the two banding patterns
20 indicate potentially differentially expressed genes.

Once potentially differentially expressed gene sequences have been identified via bulk techniques such as, for example, those described above, the differential expression of such putatively differentially expressed genes
25 should be corroborated. Corroboration may be accomplished via, for example, such well known techniques as Northern analysis and/or RT-PCR.

Upon corroboration, the differentially expressed genes may be further characterized, and may be identified as
30 target and/or fingerprint genes, as discussed, below, in Section 5.3.

Also, amplified sequences of differentially expressed genes obtained through, for example, differential display may be used to isolate full length clones of the
35 corresponding gene. The full length coding portion of the gene may readily be isolated, without undue experimentation, by molecular biological techniques well known in the art.

For example, the isolated differentially expressed amplified fragment may be labeled and used to screen a cDNA library. Alternatively, the labeled fragment may be used to screen a genomic library.

5 PCR technology may also be utilized to isolate full length cDNA sequences. As described, above, in this Section, the isolated, amplified gene fragments obtained through differential display have 5' terminal ends at some random point within the gene and have 3' terminal ends at a position
10 preferably corresponding to the 3' end of the transcribed portion of the gene. Once nucleotide sequence information from an amplified fragment is obtained, the remainder of the gene (i.e., the 5' end of the gene, when utilizing differential display) may be obtained using, for example, RT-
15 PCR.

In one embodiment of such a procedure for the identification and cloning of full length gene sequences, RNA may be isolated, following standard procedures, from an appropriate tissue or cellular source. A reverse
20 transcription reaction may then be performed on the RNA using an oligonucleotide primer complimentary to the mRNA that corresponds to the amplified fragment, for the priming of first strand synthesis. Because the primer is anti-parallel to the mRNA, extension will proceed toward the 5' end of the
25 mRNA. The resulting RNA/DNA hybrid may then be "tailed" with guanines using a standard terminal transferase reaction, the hybrid may be digested with RNAase H, and second strand synthesis may then be primed with a poly-C primer. Using the two primers, the 5' portion of the gene is amplified using
30 PCR. Sequences obtained may then be isolated and recombined with previously isolated sequences to generate a full-length cDNA of the differentially expressed genes of the invention. For a review of cloning strategies and recombinant DNA techniques, see e.g., Sambrook et al., 1989, *supra*; and
35 Ausubel et al., 1989, *supra*.

5.2. IDENTIFICATION OF PATHWAY GENES

This section describes methods for the identification of genes, termed "pathway genes", involved in cardiovascular disease. "Pathway gene", as used herein, refers to a gene whose gene product exhibits the ability to interact with gene products involved in cardiovascular disease. A pathway gene may be differentially expressed and, therefore, may additionally have the characteristics of a target and/or fingerprint gene.

- Any method suitable for detecting protein-protein interactions may be employed for identifying pathway gene products by identifying interactions between gene products and gene products known to be involved in cardiovascular disease. Such known gene products may be cellular or extracellular proteins. Those gene products which interact with such known gene products represent pathway gene products and the genes which encode them represent pathway genes.

- Among the traditional methods which may be employed are co-immunoprecipitation, crosslinking and co-purification through gradients or chromatographic columns. Utilizing procedures such as these allows for the identification of pathway gene products. Once identified, a pathway gene product may be used, in conjunction with standard techniques, to identify its corresponding pathway gene. For example, at least a portion of the amino acid sequence of the pathway gene product may be ascertained using techniques well known to those of skill in the art, such as via the Edman degradation technique (see, e.g., Creighton, 1983, Proteins: Structures and Molecular Principles, W.H. Freeman & Co., N.Y., pp.34-49). The amino acid sequence obtained may be used as a guide for the generation of oligonucleotide mixtures that can be used to screen for pathway gene sequences. Screening made be accomplished, for example by standard hybridization or PCR techniques. Techniques for the generation of oligonucleotide mixtures and screening are well-known. (See, e.g., Ausubel, *supra.*, and PCR Protocols:

A Guide to Methods and Applications, 1990, Innis, M. et al., eds. Academic Press, Inc., New York).

Additionally, methods may be employed which result in the simultaneous identification of pathway genes which
5 encode the protein interacting with a protein involved in cardiovascular disease. These methods include, for example, probing expression libraries with labeled protein known or suggested to be involved in cardiovascular disease, using this protein in a manner similar to the well known technique
10 of antibody probing of λ gt11 libraries.

One such method which detects protein interactions *in vivo*, the two-hybrid system, is described in detail for illustration only and not by way of limitation. One version of this system has been described (Chien et al., 1991, Proc.
15 Natl. Acad. Sci. USA, 88:9578-9582) and is commercially available from Clontech (Palo Alto, CA).

Briefly, utilizing such a system, plasmids are constructed that encode two hybrid proteins: one consists of the DNA-binding domain of a transcription activator protein
20 fused to a known protein, and the other consists of the activator protein's activation domain fused to an unknown protein that is encoded by a cDNA which has been recombined into this plasmid as part of a cDNA library. The plasmids are transformed into a strain of the yeast Saccharomyces
25 cerevisiae that contains a reporter gene (e.g., lacZ) whose regulatory region contains the activator's binding sites. Either hybrid protein alone cannot activate transcription of the reporter gene, the DNA-binding domain hybrid because it does not provide activation function and the activation
30 domain hybrid because it cannot localize to the activator's binding sites. Interaction of the two proteins reconstitutes the functional activator protein and results in expression of the reporter gene, which is detected by an assay for the reporter gene product.

35 The two-hybrid system or related methodology may be used to screen activation domain libraries for proteins that interact with a known "bait" gene protein. Total genomic or

5 cDNA sequences may be fused to the DNA encoding an activation domain. Such a library and a plasmid encoding a hybrid of the bait gene protein fused to the DNA-binding domain may be cotransformed into a yeast reporter strain, and the resulting transformants may be screened for those that express the reporter gene. These colonies may be purified and the library plasmids responsible for reporter gene expression may be isolated. DNA sequencing may then be used to identify the proteins encoded by the library plasmids.

10 For example, and not by way of limitation, the bait gene may be cloned into a vector such that it is translationally fused to the DNA encoding the DNA-binding domain of the GAL4 protein. Also by way of example, for the isolation of genes involved in cardiovascular disease, previously isolated genes known or suggested to play a part in cardiovascular disease may be used as the bait genes. These include but are not limited to the genes for bFGF, IGF-I, VEGF, IL-1, M-CSF, TGF β , TGF α , TNF α , HB-EGF, PDGF, IFN- γ , and GM-CSF, to name a few.

20 A cDNA library of the cell line from which proteins that interact with bait gene are to be detected can be made using methods routinely practiced in the art. According to the particular system described herein, for example, the cDNA fragments may be inserted into a vector such that they are translationally fused to the activation domain of GAL4. This library may be co-transformed along with the bait gene-GAL4 fusion plasmid into a yeast strain which contains a *lacZ* gene driven by a promoter which contains the GAL4 activation sequence. A cDNA encoded protein, fused to the GAL4 activation domain, that interacts with bait gene will reconstitute an active GAL4 protein and thereby drive expression of the *lacZ* gene. Colonies which express *lacZ* may be detected by their blue color in the presence of X-gal. The cDNA may then be purified from these strains, and used to produce and isolate the bait gene-interacting protein using techniques routinely practiced in the art.

Once a pathway gene has been identified and isolated, it may be further characterized as, for example, discussed below, in Section 5.3.

5 5.3. CHARACTERIZATION OF DIFFERENTIALLY EXPRESSED AND PATHWAY GENES

Differentially expressed genes, such as those identified via the methods discussed, above, in Section 5.1.1, pathway genes, such as those identified via the methods discussed, above, in Section 5.2, as well as genes identified by alternative means, may be further characterized by utilizing, for example, methods such as those discussed herein. Such genes will be referred to herein as "identified genes".

Analyses such as those described herein will yield information regarding the biological function of the identified genes. An assessment of the biological function of the differentially expressed genes, in addition, will allow for their designation as target and/or fingerprint genes. Specifically, any of the differentially expressed genes whose further characterization indicates that a modulation of the gene's expression or a modulation of the gene product's activity may ameliorate cardiovascular disease will be designated "target genes", as defined, above, in Section 5.1. Such target genes and target gene products, along with those discussed below, will constitute the focus of the compound discovery strategies discussed, below, in Section 5.5.

Any of the differentially expressed genes whose further characterization indicates that such modulations may not positively affect cardiovascular disease, but whose expression pattern contributes to a gene expression "fingerprint pattern" correlative of, for example, a cardiovascular disease condition will be designated a "fingerprint gene". "Fingerprint patterns" will be more fully discussed, below, in Section 5.8. It should be noted that each of the target genes may also function as

fingerprint genes, as may all or a subset of the pathway genes.

It should further be noted that the pathway genes may also be characterized according to techniques such as those described herein. Those pathway genes which yield information indicating that they are differentially expressed and that modulation of the gene's expression or a modulation of the gene product's activity may ameliorate cardiovascular disease will be also be designated "target genes". Such target genes and target gene products, along with those discussed above, will constitute the focus of the compound discovery strategies discussed, below, in Section 5.5.

It should be additionally noted that the characterization of one or more of the pathway genes may reveal a lack of differential expression, but evidence that modulation of the gene's activity or expression may, nonetheless, ameliorate cardiovascular disease symptoms. In such cases, these genes and gene products would also be considered a focus of the compound discovery strategies of Section 5.5, below.

In instances wherein a pathway gene's characterization indicates that modulation of gene expression or gene product activity may not positively affect cardiovascular disease, but whose expression is differentially expressed and which contributes to a gene expression fingerprint pattern correlative of, for example, a cardiovascular disease state, such pathway genes may additionally be designated as fingerprint genes.

Among the techniques whereby the identified genes may be further characterized, the nucleotide sequence of the identified genes, which may be obtained by utilizing standard techniques well known to those of skill in the art, may be used to further characterize such genes. For example, the sequence of the identified genes may reveal homologies to one or more known sequence motifs which may yield information regarding the biological function of the identified gene product.

Second, an analysis of the tissue distribution of the mRNA produced by the identified genes may be conducted, utilizing standard techniques well known to those of skill in the art. Such techniques may include, for example, Northern analyses and RT-PCR. Such analyses provide information as to whether the identified genes are expressed in tissues expected to contribute to cardiovascular disease. Such analyses may also provide quantitative information regarding steady state mRNA regulation, yielding data concerning which of the identified genes exhibits a high level of regulation in, preferably, tissues which may be expected to contribute to cardiovascular disease.

Such analyses may also be performed on an isolated cell population of a particular cell type derived from a given tissue. Additionally, standard in situ hybridization techniques may be utilized to provide information regarding which cells within a given tissue express the identified gene. Such analyses may provide information regarding the biological function of an identified gene relative to cardiovascular disease in instances wherein only a subset of the cells within the tissue is thought to be relevant to cardiovascular disease.

Such an in situ hybridization analysis is described in the example in Section 14, below. Specifically, the roles of the rchd502 and rchd528 genes in cardiovascular disease were further demonstrated by detecting high levels of their expression specifically within the endothelial cells of diseased tissue removed from a human cardiovascular disease patient, and not in any other cell type present in the tissue, including smooth muscle cells and macrophages. These results clearly demonstrate how detection of differentially expressed genes in the paradigms described herein leads to biologically relevant, novel, specific targets for the treatment and diagnosis of cardiovascular disease.

Third, the sequences of the identified genes may be used, utilizing standard techniques, to place the genes onto genetic maps, e.g., mouse (Copeland & Jenkins, 1991, Trends

in Genetics 7: 113-118) and human genetic maps (Cohen, et al., 1993, Nature 366: 698-701). Such mapping information may yield information regarding the genes' importance to human disease by, for example, identifying genes which map
5 near genetic regions to which known genetic cardiovascular disease tendencies map.

Fourth, the biological function of the identified genes may be more directly assessed by utilizing relevant in vivo and in vitro systems. In vivo systems may include, but
10 are not limited to, animal systems which naturally exhibit cardiovascular disease predisposition, or ones which have been engineered to exhibit such symptoms, including but not limited to the apoE-deficient atherosclerosis mouse model (Plump et al., 1992, Cell 71: 343-353). Such systems are
15 discussed in Section 5.4.4.1, below.

The use of such an in vivo system is described in detail in the example provided in Section 7, below, confirming the role of the target gene bcl-2 (see Table 1, in Section 5.4.1, below). Briefly, bcl-2 expression first was
20 shown to be down-regulated in the apoE-deficient atherosclerosis mouse model. Then, a transgenic mouse was engineered bearing the human bcl-2 gene under the control of a promoter which is induced in monocyte foam cells under atherogenic conditions. To test the effect of the induction
25 of bcl-2 under such conditions, the transgenic mouse is crossed with the apoE-deficient mouse. apoE-deficient progeny bearing the highly expressible bcl-2 gene are then examined for plaque formation and development. Reduction in plaque formation and development in these progeny confirms
30 the effectiveness of intervening in cardiovascular disease through this target gene.

In vitro systems may include, but are not limited to, cell-based systems comprising cell types known or suspected of involvement in cardiovascular disease. Such
35 systems are discussed in detail, below, in Section 5.4.4.2.

In further characterizing the biological function of the identified genes, the expression of these genes may be

modulated within the in vivo and/or in vitro systems, i.e., either over- or underexpressed, and the subsequent effect on the system then assayed. Alternatively, the activity of the product of the identified gene may be modulated by either
5 increasing or decreasing the level of activity in the in vivo and/or in vitro system of interest, and its subsequent effect then assayed.

The information obtained through such characterizations may suggest relevant methods for the
10 treatment of cardiovascular disease involving the gene of interest. For example, treatment may include a modulation of gene expression and/or gene product activity. Characterization procedures such as those described herein may indicate where such modulation should involve an increase
15 or a decrease in the expression or activity of the gene or gene product of interest.

For example, genes which are up-regulated under disease conditions may be involved in causing or exacerbating the disease condition. Treatments directed at down-
20 regulating the activity of such harmfully expressed genes will ameliorate the disease condition. On the other hand, the up-regulation of genes under disease conditions may be part of a protective response by affected cells. Treatments directed at increasing or enhancing the activity of such up-
25 regulated gene products, especially in individuals lacking normal up-regulation, will similarly ameliorate disease conditions. Such methods of treatment are discussed, below, in Section 5.6.

30 5.4. DIFFERENTIALLY EXPRESSED AND PATHWAY GENES

Identified genes, which include but are not limited to differentially expressed genes such as those identified in Section 5.1.1, above, and pathway genes, such as those identified in Section 5.2, above, are described herein.
35 Specifically, the nucleic acid sequences and gene products of such identified genes are described herein. Further, antibodies directed against the identified genes' products,

and cell- and animal-based models by which the identified genes may be further characterized and utilized are also discussed in this Section.

5 5.4.1. DIFFERENTIALLY EXPRESSED AND PATHWAY GENE SEQUENCES

 The differentially expressed and pathway genes of the invention are listed below, in Table 1. Differentially expressed and pathway gene nucleotide sequences are shown in
10 FIGS. 1-10.

 Table 1 lists differentially expressed genes identified through, for example, the paradigms discussed, above, in Section 5.1.1, and below, in the examples presented in Sections 6 through 9 and 16, below. Table 1 also
15 summarizes information regarding the further characterization of such genes.

 First, the paradigm used initially to detect the differentially expressed gene is described under the column headed "Paradigm of Original Detection". The expression
20 patterns of those genes which have been shown to be differentially expressed, for example, under one or more of the paradigm conditions described in Section 5.1.1 are summarized under the column headed "Paradigm Expression Pattern". For each of the tested genes, the paradigm which
25 was used and the difference in the expression of the gene among the samples generated is shown. "↑" indicates that gene expression is up-regulated (*i.e.*, there is an increase in the amount of detectable mRNA) among the samples generated, while "↓" indicates that gene expression is down-regulated (*i.e.*, there is a decrease in the amount of
30 detectable mRNA) among the samples generated. "Detectable" as used herein, refers to levels of mRNA which are detectable via, for example, standard Northern and/or RT-PCR techniques which are well known to those of skill in the art.

35 Cell types in which differential expression was detected are also summarized in Table 1 under the column headed "Cell Type Detected in". The column headed

"Chromosomal Location" provides the human chromosome number on which the gene is located. Additionally, in instances wherein the genes contain nucleotide sequences similar or homologous to sequences found in nucleic acid databases, 5 references to such similarities are listed.

The genes listed in Table 1 may be obtained using cloning methods well known to those skilled in the art, including but not limited to the use of appropriate probes to detect the genes within an appropriate cDNA or gDNA (genomic 10 DNA) library. (See, for example, Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratories, which is incorporated by reference herein in its entirety). Probes for the novel sequences reported herein may be obtained directly from the isolated clones 15 deposited with the NRRL or ATCC, as indicated in Table 2, below. Alternatively, oligonucleotide probes for the novel genes may be synthesized based on the DNA sequences disclosed herein in FIGS. 1-10. Such synthetic oligonucleotides may be similarly produced based on the sequences provided for the 20 previously known genes described in the following references: Cleary et al., 1986, Cell 47: 19-28 (bcl-2); Takahashi et al., 1990, J. Biochem 108: 145-148 (glutathione peroxidase); and Jones et al., 1993, J. Biol. Chem. 268: 9049-9054 (prostaglandin endoperoxide synthase II), each of which is 25 incorporated herein in its entirety.

The sequence obtained from clones containing partial coding sequences or non-coding sequences can be used to obtain the entire coding region by using the RACE method (Chenchik, et al., 1995, CLONTECHniques (X) 1: 5-8; Barnes, 30 1994, Proc. Natl. Acad. Sci. USA 91: 2216-2220; and Cheng et al., Proc. Natl. Acad. Sci. USA 91: 5695-5699). Oligonucleotides can be designed based on the sequence obtained from the partial clone that can amplify a reverse transcribed mRNA encoding the entire coding sequence. This 35 method was used, as described in the example in Section 9, below, to obtain the entire coding region of the rchd523 gene.

004060" 9824E060

Alternatively, probes can be used to screen cDNA libraries prepared from an appropriate cell or cell line in which the gene is transcribed. For example, the genes described herein that were detected in monocytes may be
5 cloned from a cDNA library prepared from monocytes isolated as described in Section 7.1.1, below. In fact, as described in detail in the example in Section 9, below, this method was applied in order to obtain the entire coding region of the rchd534 gene. Briefly, the up-regulation of this gene was
10 detected, under Paradigm D, in HUVEC's subjected to shear stress. Then, amplified partial sequence of the rchd534 gene was subcloned. The insert was then isolated and used to probe a cDNA library prepared from shear stress treated HUVEC's. A cDNA clone containing the entire rchd534 coding
15 region was detected, isolated, and sequenced.

The genes described herein that were detected in endothelial cells may also be cloned from a cDNA library constructed from endothelial cells isolated as described in Progress in Hemostasis and Thrombosis, Vol. 3, P. Spaet,
20 editor, Grune & Stratton Inc., New York, 1-28. Alternatively, the genes may be retrieved from a human placenta cDNA library (Clontech Laboratories, Palo Alto, CA), according to Takahashi et al., 1990, supra; a HUVEC cDNA library as described in Jones et al. 1993, supra; or an acute
25 lymphoblastic leukemia (SUP-B2) cDNA library as described in Cleary et al., 1986, supra, for example. Genomic DNA libraries can be prepared from any source.

30

35

TABLE 1
Differentially Expressed and Pathway Genes

Gene	Seq. ID #	Paradigm of Original Detection	Paradigm Expr. Pattern	Cell Type Detected in	Chromosomal Location	Ref	FIG.
Band 14: bel-2		B	↓	Monocytes		1	
Glutathione peroxidase		B	↓	Monocytes		2	
rchd005	SEQ ID NO: 52	C	↑	Endothelial		New 3	FIG. 1
rchd024	SEQ ID NO: 53	C	↑	Endothelial	4	New	FIG. 2
rchd032	SEQ ID NO: 54	C	↑	Endothelial		New	FIG. 3
rchd036	SEQ ID NO: 55	C	↑	Endothelial	15	New	FIG. 4
rchd502	SEQ ID NOS 56 and 57	D	↑	Endothelial		New 4	FIG. 5
rchd505: COX II		D	↑	Endothelial		5	
rchd523	SEQ ID NOS 58 and 59	D	↑	Endothelial	7	New	FIG. 6
rchd528	SEQ ID NOS 60 and 61	D	↑	Endothelial		New 6	FIG. 7
rchd530: MnSOD		D	↑	Endothelial		7	
rchd534	SEQ ID NOS 62 and 63	D	↑	Endothelial	15	New 8	FIG. 8
rchd534-long	SEQ ID NOS 64 and 65					New 8	FIG. 9
fchd540	SEQ ID NOS 66 and 67	D	↑	Endothelial	18	New 8	FIG. 10

1 Cleary et al., 1986, Cell 47: 19-28.
2 Takahashi et al., 1990, J. Biochem. 108: 145-148.

- 3 Shark Na-K-Cl cotransporter, Xu et al., 1994 Proc. Natl. Acad. Sci. U.S.A. 91: 2201-2205.
- 4 Rat matrix F/G, Hakes et al., 1991 Proc. Natl. Acad. Sci. U.S.A. 88: 6186-6190.
- 5 Jones et al., 1993, J. Biol. Chem. 268: 9049-9054.
- 6 Xenopus Xotch (homolog of *Drosophila* Notch), Coffman et al., 1990, Science 249: 1438-1441.
- 7 Heckl, 1988, Nucl. Acids Res. 16: 6224.
- 8 *Drosophila* Mothers against *dpp* (*Mad*), Sekelsky et al., 1995, Genetics 139: 1347-1358.

5

10

15

20

25

30

Table 2, below, lists isolated clones that contain sequences of the novel genes listed in Table 1. Such clones were produced from amplified sequences of the indicated differential display band which were subcloned into the TA 5 cloning vector (Invitrogen, San Diego, CA), as described in Section 6.1, below. Also listed in Table 2, below, are the strains deposited with the NRRL or ATCC which contain each such clone. Such strains were produced by transforming E. coli strain INV α F' (Invitrogen) with the indicated plasmid, 10 as described in Section 6.1, below. The names of the plasmids containing the entire coding region of a novel gene bear the prefix pFCHD, and the names of the strains carrying these plasmids bear the prefix FCHD.

15

TABLE 2

20

25

30

35

GENE		Strain Deposited	Plasmid Clone Corresponding Strain
rchd005		RCHD005	pRCHD005
rchd024		RCHD024	pRCHD024
rchd032		RCHD032	pRCHD032
rchd036		RCHD036	pRCHD036
rchd502		FCHD502SF	pFCHD502SF
		FCHD502SJ	pFCHD502SJ
		RCHD502	pRCHD502
rchd523		FCHD523	pFCHD523
		RCHD523	pRCHD523
rchd528		FCHD528A	pFCHD528A
		FCHD528B	pFCHD528B
		FCHD528C	pFCHD528C
		RCHD528	pRCHD528
rchd534	short spliceoform	FCHD534	pFCHD534
	long spliceoform		pHL6TA1A

As used herein, "differentially expressed gene" (i.e. target and fingerprint gene) or "pathway gene" refers to (a) a gene containing at least one of the DNA sequences disclosed herein (as shown in FIGS. 1-10), or contained in the clones listed in Table 2, as deposited with the ATCC or NRRL; (b) any DNA sequence that encodes the amino acid sequence encoded by the DNA sequences disclosed herein (as shown in FIGS. 1-10), contained in the clones, listed in Table 2, as deposited with the ATCC or NRRL or contained within the coding region of the gene to which the DNA sequences disclosed herein (as shown in FIGS. 1-10) or contained in the clones listed in Table 2, as deposited with the ATCC or NRRL, belong; (c) any DNA sequence that hybridizes to the complement of the coding sequences disclosed herein, contained in the clones listed in Table 2, as deposited with the ATCC or NRRL, or contained within the coding region of the gene to which the DNA sequences disclosed herein (as shown in FIGS. 1-10) or contained in the clones listed in Table 2, as deposited with the ATCC or NRRL, belong, under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at p. 2.10.3) and encodes a gene product functionally equivalent to a gene product encoded by sequences contained within the clones listed in Table 2; and/or (d) any DNA sequence that hybridizes to the complement of the coding sequences disclosed herein, (as shown in FIGS. 1-10) contained in the clones listed in Table 2, as deposited with the ATCC or NRRL or contained within the coding region of the gene to which DNA sequences disclosed herein (as shown in FIGS. 1-10) or contained in the clones, listed in Table 2, as deposited with the ATCC or NRRL, belong, under less stringent conditions, such as moderately stringent conditions, e.g., washing in 0.2xSSC/0.1% SDS at 42°C (Ausubel et al., 1989,

supra), yet which still encodes a functionally equivalent gene product.

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the DNA sequences (a) through (c), in the preceding paragraph. Such hybridization conditions may be highly stringent or less highly stringent, as described above. In instances wherein the nucleic acid molecules are deoxyoligonucleotides ("oligos"), highly stringent conditions may refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may act as target gene antisense molecules, useful, for example, in target gene regulation and/or as antisense primers in amplification reactions of target gene nucleic acid sequences. Further, such sequences may be used as part of ribozyme and/or triple helix sequences, also useful for target gene regulation. Still further, such molecules may be used as components of diagnostic methods whereby the presence of a cardiovascular disease-causing allele, may be detected.

The invention also encompasses (a) DNA vectors that contain any of the foregoing coding sequences and/or their complements (i.e., antisense); (b) DNA expression vectors that contain any of the foregoing coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences; and (c) genetically engineered host cells that contain any of the foregoing coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell. As used herein, regulatory elements include but are not limited to inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. The invention includes fragments of any of the DNA sequences disclosed herein.

In addition to the gene sequences described above,

homologues of such sequences as may, for example, be present in humans or in other species, may be identified and may be readily isolated, without undue experimentation, by molecular biological techniques well known in the art. Further, there
5 may exist genes at other genetic loci within the genome that encode proteins which have extensive homology to one or more domains of such gene products. These genes may also be identified via similar techniques.

For example, the isolated differentially expressed
10 gene sequence may be labeled and used to screen a cDNA library constructed from mRNA obtained from the organism of interest. Hybridization conditions will be of a lower stringency when the cDNA library was derived from an organism different from the type of organism from which the labeled
15 sequence was derived. Alternatively, the labeled fragment may be used to screen a genomic library derived from the organism of interest, again, using appropriately stringent conditions. Such low stringency conditions will be well known to those of skill in the art, and will vary predictably
20 depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, Cold Springs Harbor Press, N.Y.; and Ausubel et al., 1989, Current
25 Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y.

Further, a previously unknown differentially expressed or pathway gene-type sequence may be isolated by performing PCR using two degenerate oligonucleotide primer
30 pools designed on the basis of amino acid sequences within the gene of interest. The template for the reaction may be cDNA obtained by reverse transcription of mRNA prepared from human or non-human cell lines or tissue known or suspected to express a differentially expressed or pathway gene allele.

35 The PCR product may be subcloned and sequenced to insure that the amplified sequences represent the sequences of a differentially expressed or pathway gene-like nucleic

acid sequence. The PCR fragment may then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment may be labeled and used to screen a bacteriophage cDNA library. Alternatively, the
5 labeled fragment may be used to screen a genomic library.

PCR technology may also be utilized to isolate full length cDNA sequences. For example, RNA may be isolated, following standard procedures, from an appropriate cellular or tissue source. A reverse transcription reaction may be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" with guanines using a standard terminal transferase reaction, the hybrid may be digested with RNAase H, and second strand synthesis may then be primed with a poly-C primer. Thus, cDNA sequences upstream of the amplified fragment may easily be isolated. For a review of cloning strategies which may be used, see e.g., Sambrook et al., 1989, *supra*.

20 In cases where the differentially expressed or
pathway gene identified is the normal, or wild type, gene,
this gene may be used to isolate mutant alleles of the gene.
Such an isolation is preferable in processes and disorders
which are known or suspected to have a genetic basis. Mutant
25 alleles may be isolated from individuals either known or
suspected to have a genotype which contributes to
cardiovascular disease symptoms. Mutant alleles and mutant
allele products may then be utilized in the therapeutic and
diagnostic assay systems described below.

30 A cDNA of the mutant gene may be isolated, for example, by using PCR, a technique which is well known to those of skill in the art. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or
35 suspected to be expressed in an individual putatively carrying the mutant allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is

then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR, cloned into a suitable vector, and subjected to DNA sequence

5 analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant gene to that of the normal gene, the mutation(s) responsible for the loss or alteration of function of the mutant gene product can be ascertained.

10 Alternatively, a genomic or cDNA library can be constructed and screened using DNA or RNA, respectively, from a tissue known to or suspected of expressing the gene of interest in an individual suspected of or known to carry the mutant allele. The normal gene or any suitable fragment
15 thereof may then be labeled and used as a probed to identify the corresponding mutant allele in the library. The clone containing this gene may then be purified through methods routinely practiced in the art, and subjected to sequence analysis as described, above, in this Section.

20 Additionally, an expression library can be constructed utilizing DNA isolated from or cDNA synthesized from a tissue known to or suspected of expressing the gene of interest in an individual suspected of or known to carry the mutant allele. In this manner, gene products made by the
25 putatively mutant tissue may be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against the normal gene product, as described, below, in Section 5.4.3. (For screening techniques, see, for example, Harlow, E. and Lane, eds.,
30 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.) In cases where the mutation results in an expressed gene product with altered function (e.g., as a result of a missense mutation), a polyclonal set of antibodies are likely to cross-react with the mutant gene
35 product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis as described in this Section, above.

5.4.2. DIFFERENTIALLY EXPRESSED AND PATHWAY GENE PRODUCTS

Differentially expressed and pathway gene products include those proteins encoded by the differentially expressed and pathway gene sequences described in Section 5.4.1, above. Specifically, differentially expressed and pathway gene products may include differentially expressed and pathway gene polypeptides encoded by the differentially expressed and pathway gene sequences contained in the clones listed in Table 2, above, as deposited with the NRRL or ATCC, or contained in the coding regions of the genes to which DNA sequences disclosed herein (in FIGS. 1-10) or contained in the clones, listed in Table 2, as deposited with the NRRL or ATCC, belong, for example.

In addition, differentially expressed and pathway gene products may include proteins that represent functionally equivalent gene products. Such an equivalent differentially expressed or pathway gene product may contain deletions, additions or substitutions of amino acid residues within the amino acid sequence encoded by the differentially expressed or pathway gene sequences described, above, in Section 5.4.1, but which result in a silent change, thus producing a functionally equivalent differentially expressed on pathway gene product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved.

For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid. "Functionally equivalent", as utilized herein, refers to a protein capable of exhibiting a substantially similar in vivo activity as the endogenous differentially expressed or pathway gene products encoded by

the differentially expressed or pathway gene sequences described in Section 5.4.1, above. Alternatively, when utilized as part of assays such as those described, below, in Section 5.5, "functionally equivalent" may refer to peptides
5 capable of interacting with other cellular or extracellular molecules in a manner substantially similar to the way in which the corresponding portion of the endogenous differentially expressed or pathway gene product would.

The differentially expressed or pathway gene
10 products may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing the differentially expressed or pathway gene polypeptides and peptides of the invention by expressing nucleic acid encoding differentially expressed or pathway
15 gene sequences are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing differentially expressed or pathway gene protein coding sequences and appropriate transcriptional/translational control signals. These methods
20 include, for example, in vitro recombinant DNA techniques, synthetic techniques and in vivo recombination/genetic recombination. See, for example, the techniques described in Sambrook et al., 1989, *supra*, and Ausubel et al., 1989, *supra*. Alternatively, RNA capable of encoding differentially
25 expressed or pathway gene protein sequences may be chemically synthesized using, for example, synthesizers. See, for example, the techniques described in "Oligonucleotide Synthesis", 1984, Gait, M.J. ed., IRL Press, Oxford, which is incorporated by reference herein in its entirety.

30 A variety of host-expression vector systems may be utilized to express the differentially expressed or pathway gene coding sequences of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also
35 represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, exhibit the differentially expressed or pathway gene protein of the

invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing differentially
5 expressed or pathway gene protein coding sequences; yeast (e.g. *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing the differentially expressed or pathway gene protein coding sequences; insect cell systems infected with recombinant virus expression
10 vectors (e.g., baculovirus) containing the differentially expressed or pathway gene protein coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expres-
15 sion vectors (e.g., Ti plasmid) containing differentially expressed or pathway gene protein coding sequences; or mammalian cell systems (e.g. COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g.,
20 metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use
25 intended for the differentially expressed or pathway gene protein being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of antibodies or to screen peptide libraries, for example, vectors which direct the expression of high levels of fusion
30 protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which the differentially expressed or pathway gene protein coding sequence may be ligated individually into
35 the vector in frame with the *lac Z* coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster,

1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be
5 purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene protein can be released from the GST
10 moiety.

In a preferred embodiment, full length cDNA sequences are appended with in-frame Bam HI sites at the amino terminus and Eco RI sites at the carboxyl terminus using standard PCR methodologies (Innis et al., 1990, *supra*)
15 and ligated into the pGEX-2TK vector (Pharmacia, Uppsala, Sweden). The resulting cDNA construct contains a kinase recognition site at the amino terminus for radioactive labelling and glutathione S-transferase sequences at the carboxyl terminus for affinity purification (Nilsson, et al.,
20 1985, EMBO J. 4: 1075; Zabeau and Stanley, 1982, EMBO J. 1: 1217.

In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda*
25 cells. The differentially expressed or pathway gene coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of differentially
30 expressed or pathway gene coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera*
35 *frugiperda* cells in which the inserted gene is expressed. (E.g., see Smith et al., 1983, J. Virol. 46: 584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the differentially expressed or pathway gene coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing differentially expressed or pathway gene protein in infected hosts. (E.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted differentially expressed or pathway gene coding sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire differentially expressed or pathway gene, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of the differentially expressed or pathway gene coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al., 1987, Methods in Enzymol. 153:516-544).

In a preferred embodiment, cDNA sequences encoding the full-length open reading frames are ligated into pCMV β replacing the β -galactosidase gene such that cDNA expression

is driven by the CMV promoter (Alam, 1990, Anal. Biochem. 188: 245-254; MacGregor & Caskey, 1989, Nucl. Acids Res. 17: 2365; Norton & Corrin, 1985, Mol. Cell. Biol. 5: 281).

In addition, a host cell strain may be chosen which
5 modulates the expression of the inserted sequences, or
modifies and processes the gene product in the specific
fashion desired. Such modifications (e.g., glycosylation)
and processing (e.g., cleavage) of protein products may be
important for the function of the protein. Different host
10 cells have characteristic and specific mechanisms for the
post-translational processing and modification of proteins.
Appropriate cell lines or host systems can be chosen to
ensure the correct modification and processing of the foreign
protein expressed. To this end, eukaryotic host cells which
15 possess the cellular machinery for proper processing of the
primary transcript, glycosylation, and phosphorylation of the
gene product may be used. Such mammalian host cells include
but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293,
3T3, WI38, etc.

20 For long-term, high-yield production of recombinant
proteins, stable expression is preferred. For example, cell
lines which stably express the differentially expressed or
pathway gene protein may be engineered. Rather than using
expression vectors which contain viral origins of
25 replication, host cells can be transformed with DNA
controlled by appropriate expression control elements (e.g.,
promoter, enhancer, sequences, transcription terminators,
polyadenylation sites, etc.), and a selectable marker.
Following the introduction of the foreign DNA, engineered
30 cells may be allowed to grow for 1-2 days in an enriched
media, and then are switched to a selective media. The
selectable marker in the recombinant plasmid confers
resistance to the selection and allows cells to stably
integrate the plasmid into their chromosomes and grow to form
35 foci which in turn can be cloned and expanded into cell
lines. This method may advantageously be used to engineer
cell lines which express the differentially expressed or

pathway gene protein. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the differentially expressed or pathway gene protein.

- 5 A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and
- 10 adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes can be employed in tk⁻, hgp^rt⁻ or ap^rt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA
- 15 77:3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150:1); and
- 20 hyg^r, which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30:147) genes.

- An alternative fusion protein system allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl.
- 25 Acad. Sci. USA 88: 8972-8976). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant
- 30 vaccinia virus are loaded onto Ni²⁺·nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

- When used as a component in assay systems such as those described, below, in Section 5.5, the differentially
- 35 expressed or pathway gene protein may be labeled, either directly or indirectly, to facilitate detection of a complex formed between the differentially expressed or pathway gene

protein and a test substance. Any of a variety of suitable labeling systems may be used including but not limited to radioisotopes such as ¹²⁵I; enzyme labelling systems that generate a detectable colorimetric signal or light when
5 exposed to substrate; and fluorescent labels.

Where recombinant DNA technology is used to produce the differentially expressed or pathway gene protein for such assay systems, it may be advantageous to engineer fusion proteins that can facilitate labeling, immobilization and/or
10 detection.

Indirect labeling involves the use of a protein, such as a labeled antibody, which specifically binds to either a differentially expressed or pathway gene product. Such antibodies include but are not limited to polyclonal,
15 monoclonal, chimeric, single chain, Fab fragments and fragments produced by an Fab expression library.

5.4.3. DIFFERENTIALLY EXPRESSED OR PATHWAY GENE PRODUCT ANTIBODIES

20 Described herein are methods for the production of antibodies capable of specifically recognizing one or more differentially expressed or pathway gene epitopes. Such antibodies may include, but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or
25 chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Such antibodies may be used, for example, in the detection of a fingerprint,
30 target, or pathway gene in a biological sample, or, alternatively, as a method for the inhibition of abnormal target gene activity. Thus, such antibodies may be utilized as part of cardiovascular disease treatment methods, and/or may be used as part of diagnostic techniques whereby patients
35 may be tested for abnormal levels of fingerprint, target, or pathway gene proteins, or for the presence of abnormal forms of the such proteins.

For the production of antibodies to a differentially expressed or pathway gene, various host animals may be immunized by injection with a differentially expressed or pathway gene protein, or a portion thereof.

- 5 Such host animals may include but are not limited to rabbits, mice, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide,
- 10 surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*.
- 15 In a preferred embodiment, peptide sequences corresponding to amino sequences of target gene products were selected and submitted to Research Genetics (Huntsville, AL) for synthesis and antibody production. Peptides were modified as described (Tam, J.P., 1988, Proc. Natl. Acad.
- 20 Sci. USA 85: 5409-5413; Tam, J.P., and Zavala, F., 1989, J. Immunol. Methods 124: 53-61; Tam, J.P., and Lu, Y.A., 1989, Proc. Natl. Acad. Sci. USA 86: 9084-9088), emulsified in an equal volume of Freund's adjuvant and injected into rabbits at 3 to 4 subcutaneous dorsal sites for a total volume of 1.0
- 25 ml (0.5 mg peptide) per immunization. The animals were boosted after 2 and 6 weeks and bled at weeks 4, 8, and 10. The blood was allowed to clot and serum was collected by centrifugation.

- Polyclonal antibodies are heterogeneous populations
- 30 of antibody molecules derived from the sera of animals immunized with an antigen, such as target gene product, or an antigenic functional derivative thereof. For the production of polyclonal antibodies, host animals such as those described above, may be immunized by injection with
 - 35 differentially expressed or pathway gene product supplemented with adjuvants as also described above.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, may be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture.

- 5 These include, but are not limited to the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-
10 hybridoma technique (Cole et al., 1985, Monoclonal Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated in
15 vitro or in vivo. Production of high titers of mAbs in vivo makes this the presently preferred method of production.

- In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, Proc. Natl. Acad. Sci., 81:6851-6855; Neuberger et al., 1984,
20 Nature, 312:604-608; Takeda et al., 1985, Nature, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which
25 different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region.

- Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778;
30 Bird, 1988, Science 242:423-426; Huston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward et al., 1989, Nature 334:544-546) can be adapted to produce differentially expressed or pathway gene-single chain antibodies. Single chain antibodies are formed by linking the heavy and light
35 chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

5.4.4. CELL- AND ANIMAL-BASED MODEL SYSTEMS

Described herein are cell- and animal-based systems which act as models for cardiovascular disease. These systems may be used in a variety of applications. For example, the cell- and animal-based model systems may be used to further characterize differentially expressed and pathway genes, as described, above, in Section 5.3. Such further characterization may, for example, indicate that a differentially expressed gene is a target gene. Second, such assays may be utilized as part of screening strategies designed to identify compounds which are capable of ameliorating cardiovascular disease symptoms, as described, below, in Section 5.5.4. Thus, the animal- and cell-based models may be used to identify drugs, pharmaceuticals, therapies and interventions which may be effective in treating cardiovascular disease. In addition, as described in detail, below, in Section 5.7.1, such animal models may be used to determine the LD₅₀ and the ED₅₀ in animal subjects, and such data can be used to determine the in vivo efficacy of potential cardiovascular disease treatments.

5.4.4.1. ANIMAL-BASED SYSTEMS

Animal-based model systems of cardiovascular disease may include, but are not limited to, non-recombinant and engineered transgenic animals.

Non-recombinant animal models for cardiovascular disease may include, for example, genetic models. Such genetic cardiovascular disease models may include, for example, apoB or apoR deficient pigs (Rapacz, et al., 1986, 5 Science 234:1573-1577) and Watanabe heritable hyperlipidemic (WHHL) rabbits (Kita et al., 1987, Proc. Natl. Acad. Sci USA 84: 5928-5931).

Non-recombinant, non-genetic animal models of atherosclerosis may include, for example, pig, rabbit, or rat 10 models in which the animal has been exposed to either chemical wounding through dietary supplementation of LDL, or mechanical wounding through balloon catheter angioplasty, for example.

Additionally, animal models exhibiting 15 cardiovascular disease symptoms may be engineered by utilizing, for example, target gene sequences such as those described, above, in Section 5.4.1, in conjunction with techniques for producing transgenic animals that are well known to those of skill in the art. For example, target gene 20 sequences may be introduced into, and overexpressed in, the genome of the animal of interest, or, if endogenous target gene sequences are present, they may either be overexpressed or, alternatively, be disrupted in order to underexpress or inactivate target gene expression, such as described for the 25 disruption of apoE in mice (Plump et al., 1992, Cell 71: 343-353).

In order to overexpress a target gene sequence, the coding portion of the target gene sequence may be ligated to a regulatory sequence which is capable of driving gene 30 expression in the animal and cell type of interest. Such regulatory regions will be well known to those of skill in the art, and may be utilized in the absence of undue experimentation.

The use of such a genetically engineered animal- 35 based system is described in detail in the example provided in Section 7, below, for the target gene bcl-2 (see Table 1, in Section 5.4.1, above). Briefly, bcl-2 expression first

was shown to be down-regulated in the apoE-deficient atherosclerosis mouse model. Then, a transgenic mouse was engineered bearing the human bcl-2 gene under the control of a promoter which is induced under atherogenic conditions. To
5 test the effect of the induction of bcl-2 under such conditions, the transgenic mouse is crossed with the apoE-deficient mouse. apoE-deficient progeny bearing the highly expressible bcl-2 gene are then examined for plaque formation and development. Reduction in plaque formation and
10 development in these progeny confirms the effectiveness of intervening in cardiovascular disease through this target gene.

For underexpression of an endogenous target gene sequence, such a sequence may be isolated and engineered such
15 that when reintroduced into the genome of the animal of interest, the endogenous target gene alleles will be inactivated. Preferably, the engineered target gene sequence is introduced via gene targeting such that the endogenous target sequence is disrupted upon integration of the
20 engineered target gene sequence into the animal's genome. Gene targeting is discussed, below, in this Section.

Animals of any species, including, but not limited to, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, goats, and non-human primates, e.g., baboons, monkeys, and
25 chimpanzees may be used to generate cardiovascular disease animal models.

Any technique known in the art may be used to introduce a target gene transgene into animals to produce the founder lines of transgenic animals. Such techniques
30 include, but are not limited to pronuclear microinjection (Hoppe, P.C. and Wagner, T.E., 1989, U.S. Pat. No. 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten et al., 1985, Proc. Natl. Acad. Sci., USA 82:6148-6152); gene targeting in embryonic stem cells
35 (Thompson et al., 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer (Lavitrano et al., 1989, Cell 57:717-

723); etc. For a review of such techniques, see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.

The present invention provides for transgenic
5 animals that carry the transgene in all their cells, as well
as animals which carry the transgene in some, but not all
their cells, i.e., mosaic animals. The transgene may be
integrated as a single transgene or in concatamers, e.g.,
head-to-head tandems or head-to-tail tandems. The transgene
10 may also be selectively introduced into and activated in a
particular cell type by following, for example, the teaching
of Lasko et al. (Lasko, M. et al., 1992, Proc. Natl. Acad.
Sci. USA 89: 6232-6236). The regulatory sequences required
for such a cell-type specific activation will depend upon the
15 particular cell type of interest, and will be apparent to
those of skill in the art. When it is desired that the
target gene transgene be integrated into the chromosomal site
of the endogenous target gene, gene targeting is preferred.
Briefly, when such a technique is to be utilized, vectors
20 containing some nucleotide sequences homologous to the
endogenous target gene of interest are designed for the
purpose of integrating, via homologous recombination with
chromosomal sequences, into and disrupting the function of
the nucleotide sequence of the endogenous target gene. The
25 transgene may also be selectively introduced into a
particular cell type, thus inactivating the endogenous gene
of interest in only that cell type, by following, for
example, the teaching of Gu et al. (Gu, et al., 1994, Science
265: 103-106). The regulatory sequences required for such a
30 cell-type specific inactivation will depend upon the
particular cell type of interest, and will be apparent to
those of skill in the art. Recombinant methods for
expressing target genes are described in Section 5.4.2,
above.

35 Once transgenic animals have been generated, the
expression of the recombinant target gene and protein may be
assayed utilizing standard techniques. Initial screening may

be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may
5 also be assessed using techniques which include but are not limited to Northern blot analysis of tissue samples obtained from the animal, in situ hybridization analysis, and RT-PCR. Samples of target gene-expressing tissue, may also be evaluated immunocytochemically using antibodies specific for
10 the target gene transgene gene product of interest.

The target gene transgenic animals that express target gene mRNA or target gene transgene peptide (detected immunocytochemically, using antibodies directed against the target gene product's epitopes) at easily detectable levels should then be further evaluated to identify those animals which display characteristic cardiovascular disease symptoms. Such symptoms may include, for example, increased prevalence and size of fatty streaks and/or cardiovascular disease plaques.

Once target gene transgenic founder animals are produced, they may be bred, inbred, outbred, or crossbred to

produce colonies of the particular animal. Examples of such breeding strategies include but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of
5 separate lines in order to produce compound target gene transgenics that express the target gene transgene of interest at higher levels because of the effects of additive expression of each target gene transgene; crossing of heterozygous transgenic animals to produce animals homozygous
10 for a given integration site in order both to augment expression and eliminate the possible need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; breeding animals to different inbred genetic backgrounds so
15 as to examine effects of modifying alleles on expression of the target gene transgene and the development of cardiovascular disease symptoms. One such approach is to cross the target gene transgenic founder animals with a wild type strain to produce an F1 generation that exhibits
20 cardiovascular disease symptoms. The F1 generation may then be inbred in order to develop a homozygous line, if it is found that homozygous target gene transgenic animals are viable.

25 5.4.4.2. CELL-BASED ASSAYS

Cells that contain and express target gene sequences which encode target gene protein, and, further, exhibit cellular phenotypes associated with cardiovascular disease, may be utilized to identify compounds that exhibit
30 anti-cardiovascular disease activity.

Such cells may include non-recombinant monocyte cell lines, such as U937 (ATCC# CRL-1593), THP-1 (ATCC# TIB-202), and P388D1 (ATCC# TIB-63); endothelial cells such as HUVEC's and bovine aortic endothelial cells (BAEC's); as well
35 as generic mammalian cell lines such as HeLa cells and COS cells, e.g., COS-7 (ATCC# CRL-1651). Further, such cells may include recombinant, transgenic cell lines. For example, the

cardiovascular disease animal models of the invention, discussed, above, in Section 5.4.4.1, may be used to generate cell lines, containing one or more cell types involved in cardiovascular disease, that can be used as cell culture models for this disorder. While primary cultures derived from the cardiovascular disease transgenic animals of the invention may be utilized, the generation of continuous cell lines is preferred. For examples of techniques which may be used to derive a continuous cell line from the transgenic animals, see Small et al., 1985, Mol. Cell Biol. 5:642-648.

Alternatively, cells of a cell type known to be involved in cardiovascular disease may be transfected with sequences capable of increasing or decreasing the amount of target gene expression within the cell. For example, target gene sequences may be introduced into, and overexpressed in, the genome of the cell of interest, or, if endogenous target gene sequences are present, they may be either overexpressed or, alternatively disrupted in order to underexpress or inactivate target gene expression.

In order to overexpress a target gene sequence, the coding portion of the target gene sequence may be ligated to a regulatory sequence which is capable of driving gene expression in the cell type of interest. Such regulatory regions will be well known to those of skill in the art, and may be utilized in the absence of undue experimentation. Recombinant methods for expressing target genes are described in Section 5.4.2, above.

For underexpression of an endogenous target gene sequence, such a sequence may be isolated and engineered such that when reintroduced into the genome of the cell type of interest, the endogenous target gene alleles will be inactivated. Preferably, the engineered target gene sequence is introduced via gene targeting such that the endogenous target sequence is disrupted upon integration of the engineered target gene sequence into the cell's genome. Transfection of host cells with target genes is discussed, above, in Section 5.4.4.1.

Cells treated with compounds or transfected with target genes can be examined for phenotypes associated with cardiovascular disease. In the case of monocytes, such phenotypes include but are not limited to increases in rates of LDL uptake, adhesion to endothelial cells, transmigration, foam cell formation, fatty streak formation, and production by foam cells of growth factors such as bFGF, IGF-I, VEGF, IL-1, M-CSF, TGF β , TGF α , TNF α , HB-EGF, PDGF, IFN- γ , and GM-CSF. Transmigration rates, for example, may be measured using the in vitro system of Navab et al., described in Section 5.1.1.3, above, by quantifying the number of monocytes that migrate across the endothelial monolayer and into the collagen layer of the subendothelial space.

Similarly, HUVEC's can be treated with test compounds or transfected with genetically engineered target genes described in Section 5.4.2, above. The HUVEC's can then be examined for phenotypes associated with cardiovascular disease, including, but not limited to changes in cellular morphology, cell proliferation, cell migration, and mononuclear cell adhesion; or for the effects on production of other proteins involved in cardiovascular disease such as ICAM, VCAM, PDGF- β , and E-selectin.

Transfection of target gene sequence nucleic acid may be accomplished by utilizing standard techniques. See, for example, Ausubel, 1989, *supra*. Transfected cells should be evaluated for the presence of the recombinant target gene sequences, for expression and accumulation of target gene mRNA, and for the presence of recombinant target gene protein production. In instances wherein a decrease in target gene expression is desired, standard techniques may be used to demonstrate whether a decrease in endogenous target gene expression and/or in target gene product production is achieved.

5.5. SCREENING ASSAYS FOR COMPOUNDS THAT INTERACT
WITH THE TARGET GENE PRODUCT AND/OR
MODULATE TARGET GENE EXPRESSION

5 The following assays are designed to identify
compounds that bind to target gene products, bind to other
cellular or extracellular proteins that interact with a
target gene product, and interfere with the interaction of
the target gene product with other cellular or extracellular
proteins. Such compounds can act as the basis for
amelioration of such cardiovascular diseases as
10 atherosclerosis, ischemia/reperfusion, hypertension,
restenosis, and arterial inflammation by modulating the
activity of the protein products of target genes. Such
compounds may also act as the basis for the amelioration of
15 fibroproliferative and oncogenic related disorders, including
tumorigenesis and the vascularization of tumors. Such
compounds may include, but are not limited to peptides,
antibodies, or small organic or inorganic compounds. Methods
for the identification of such compounds are described in
20 Section 5.5.1, below. Such compounds may also include other
cellular proteins. Methods for the identification of such
cellular proteins are described, below, in Section 5.5.2.

For example, such assays can be used to identify
compounds that bind to the rchd534 protein, the rchd534-long
25 protein, or the fchd540 protein, including compounds that
interfere with the interaction of the the rchd534 protein
with itself, the rchd534-long protein with itself, or the
fchd540 protein with either the rchd534 protein or the
rchd534-long protein.

30 Compounds identified via assays such as those
described herein may be useful, for example, in elaborating
the biological function of the target gene product, and for
ameliorating cardiovascular disease. In instances whereby a
cardiovascular disease condition results from an overall
35 lower level of target gene expression and/or target gene
product in a cell or tissue, compounds that interact with the
target gene product may include compounds which accentuate or
amplify the activity of the bound target gene protein. Such

compounds would bring about an effective increase in the level of target gene product activity, thus ameliorating symptoms.

In some cases, a target gene observed to be up-regulated under disease conditions may be exerting a protective effect. Compounds that enhance the expression of such up-regulated genes, or the activity of their gene products, would also ameliorate disease symptoms, especially in individuals whose target gene is not normally up-regulated.

In other instances mutations within the target gene may cause aberrant types or excessive amounts of target gene proteins to be made which have a deleterious effect that leads to cardiovascular disease. Similarly, physiological conditions may cause an excessive increase in target gene expression leading to cardiovascular disease. In such cases, compounds that bind target gene protein may be identified that inhibit the activity of the bound target gene protein. Assays for testing the effectiveness of compounds, identified by, for example, techniques such as those described in this Section are discussed, below, in Section 5.5.4.

5.5.1. IN VITRO SCREENING ASSAYS FOR COMPOUNDS THAT BIND TO THE TARGET GENE PRODUCT

In vitro systems may be designed to identify compounds capable of binding the target gene of the invention. Such compounds may include, but are not limited to, peptides made of D-and/or L-configuration amino acids (in, for example, the form of random peptide libraries; see e.g., Lam, K.S. et al., 1991, Nature 354:82-84), phosphopeptides (in, for example, the form of random or partially degenerate, directed phosphopeptide libraries; see, e.g., Songyang, Z. et al., 1993, Cell 72:767-778), antibodies, and small organic or inorganic molecules. Compounds identified may be useful, for example, in modulating the activity of target gene proteins, preferably mutant target gene proteins, may be useful in elaborating the

biological function of the target gene protein, may be utilized in screens for identifying compounds that disrupt normal target gene interactions, or may in themselves disrupt such interactions.

- 5 The principle of the assays used to identify compounds that bind to the target gene protein involves preparing a reaction mixture of the target gene protein and the test compound under conditions and for a time sufficient to allow the two components to interact and bind, thus
- 10 forming a complex which can be removed and/or detected in the reaction mixture. These assays can be conducted in a variety of ways. For example, one method to conduct such an assay would involve anchoring the target gene or the test substance onto a solid phase and detecting target gene/test substance
- 15 complexes anchored on the solid phase at the end of the reaction. In one embodiment of such a method, the target gene protein may be anchored onto a solid surface, and the test compound, which is not anchored, may be labeled, either directly or indirectly.
- 20 In practice, microtitre plates are conveniently utilized. The anchored component may be immobilized by non-covalent or covalent attachments. Non-covalent attachment may be accomplished simply by coating the solid surface with a solution of the protein and drying. Alternatively, an
- 25 immobilized antibody, preferably a monoclonal antibody, specific for the protein may be used to anchor the protein to the solid surface. The surfaces may be prepared in advance and stored.

- In order to conduct the assay, the nonimmobilized
- 30 component is added to the coated surface containing the anchored component. After the reaction is complete, unreacted components are removed (e.g., by washing) under conditions such that any complexes formed will remain immobilized on the solid surface. The detection of complexes
- 35 anchored on the solid surface can be accomplished in a number of ways. Where the previously nonimmobilized component is pre-labeled, the detection of label immobilized on the

surface indicates that complexes were formed. Where the previously nonimmobilized component is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the
5 previously nonimmobilized component (the antibody, in turn, may be directly labeled or indirectly labeled with a labeled anti-Ig antibody).

Alternatively, a reaction can be conducted in a liquid phase, the reaction products separated from unreacted
10 components, and complexes detected; e.g., using an immobilized antibody specific for target gene product or the test compound to anchor any complexes formed in solution, and a labeled antibody specific for the other component of the possible complex to detect anchored complexes.

15 Compounds that are shown to bind to a particular target gene product through one of the methods described above can be further tested for their ability to elicit a biochemical response from the target gene protein. A particular embodiment is described herein for receptor
20 proteins involved in signal transduction, including but not limited to the rchd523 gene product. Compounds that interact with a target gene product receptor domain, can be screened for their ability to function as ligands, i.e., to bind to the receptor protein in a manner that triggers the signal
25 transduction pathway. Useful receptor fragments or analogs in the invention are those which interact with ligand. The receptor component can be assayed functionally, i.e., for its ability to bind ligand and mobilize Ca^{++} (see below). These assays include, as components, ligand and a recombinant
30 target gene product (or a suitable fragment or analog) configured to permit detection of binding.

For example, and not by way of limitation, a recombinant receptor may be used to screen for ligands by its ability to mediate ligand-dependent mobilization of calcium.
35 Cells, preferably myeloma cells or *Xenopus* oocytes, transfected with a target gene expression vector (constructed according to the methods described in Section 5.4.2, above)

are loaded with FURA-2 or INDO-1 by standard techniques. Mobilization of Ca^{2+} induced by ligand is measured by fluorescence spectroscopy as previously described (Grynkiewicz et al., 1985, *J. Biol. Chem.* 260:3440). Ligands that react with the target gene product receptor domain, therefore, can be identified by their ability to produce a fluorescent signal. Their receptor binding activities can be quantified and compared by measuring the level of fluorescence produced over background.

10 The rchd523 gene product consists of a G protein-coupled receptor with multiple transmembrane domains. The Ca^{2+} mobilization assay, therefore, can be used to screen compounds that are ligands of the rchd523 receptor. This screening method is described in detail with respect to
15 rchd523 in the example in Section 12, below. Identification of rchd523 ligand, and measuring the activity of the ligand-receptor complex, leads to the identification of antagonists of this interaction, as described in Section 5.5.3, below. Such antagonists are useful in the treatment of
20 cardiovascular disease.

5.5.2. ASSAYS FOR CELLULAR OR EXTRACELLULAR PROTEINS THAT INTERACT WITH THE TARGET GENE PRODUCT

25 Any method suitable for detecting protein-protein interactions may be employed for identifying novel target protein-cellular or extracellular protein interactions. These methods are outlined in Section 5.2., *supra*, for the identification of pathway genes, and may be utilized herein
30 with respect to the identification of proteins which interact with identified target proteins. In such a case, the target gene serves as the known "bait" gene.

35 5.5.3. ASSAYS FOR COMPOUNDS THAT INTERFERE WITH INTERACTION BETWEEN TARGET GENE PRODUCT AND OTHER COMPOUNDS

 The target gene proteins of the invention may, in vivo, interact with one or more cellular or extracellular

030498 9824E050

proteins. Such proteins may include, but are not limited to, those proteins identified via methods such as those described, above, in Section 5.5.2. For the purposes of this discussion, target gene products and such cellular and
5 extracellular proteins are referred to herein as "binding partners". Compounds that disrupt such interactions may be useful in regulating the activity of the target gene proteins, especially mutant target gene proteins. Such compounds may include, but are not limited to molecules such
10 as antibodies, peptides, and the like described in Section 5.5.1. above.

The basic principle of the assay systems used to identify compounds that interfere with the interaction between the target gene protein, and its cellular or
15 extracellular protein binding partner or partners involves preparing a reaction mixture containing the target gene protein and the binding partner under conditions and for a time sufficient to allow the two proteins to interact and bind, thus forming a complex. In order to test a compound
20 for inhibitory activity, the reaction mixture is prepared in the presence and absence of the test compound. The test compound may be initially included in the reaction mixture or may be added at a time subsequent to the addition of target gene and its cellular or extracellular binding partner.
25 Control reaction mixtures are incubated without the test compound or with a placebo. The formation of any complexes between the target gene protein and the cellular or extracellular binding partner is then detected. The formation of a complex in the control reaction, but not in
30 the reaction mixture containing the test compound, indicates that the compound interferes with the interaction of the target gene protein and the interactive binding partner protein. Additionally, complex formation within reaction mixtures containing the test compound and a normal target
35 gene protein may also be compared to complex formation within reaction mixtures containing the test compound and mutant target gene protein. This comparison may be important in

those cases wherein it is desirable to identify compounds that disrupt interactions of mutant but not normal target gene proteins.

The assay for compounds that interfere with the interaction of the binding partners can be conducted in a heterogeneous or homogeneous format. Heterogeneous assays involve anchoring one of the binding partners onto a solid phase and detecting complexes anchored on the solid phase at the end of the reaction. In homogeneous assays, the entire reaction is carried out in a liquid phase. In either approach, the order of addition of reactants can be varied to obtain different information about the compounds being tested. For example, test compounds that interfere with the interaction between the binding partners, e.g., by competition, can be identified by conducting the reaction in the presence of the test substance; i.e., by adding the test substance to the reaction mixture prior to or simultaneously with the target gene protein and interactive cellular or extracellular protein. Alternatively, test compounds that disrupt preformed complexes, e.g. compounds with higher binding constants that displace one of the binding partners from the complex, can be tested by adding the test compound to the reaction mixture after complexes have been formed. The various formats are described briefly below.

In a heterogeneous assay system, either the target gene protein or the interactive cellular or extracellular binding partner protein, is anchored onto a solid surface, and its binding partner, which is not anchored, is labeled, either directly or indirectly. In practice, microtitre plates are conveniently utilized. The anchored species may be immobilized by non-covalent or covalent attachments. Non-covalent attachment may be accomplished simply by coating the solid surface with a solution of the protein and drying. Alternatively, an immobilized antibody specific for the protein may be used to anchor the protein to the solid surface. The surfaces may be prepared in advance and stored.

864030-9824E050

In order to conduct the assay, the binding partner of the immobilized species is exposed to the coated surface with or without the test compound. After the reaction is complete, unreacted components are removed (e.g., by washing) and any complexes formed will remain immobilized on the solid surface. The detection of complexes anchored on the solid surface can be accomplished in a number of ways. Where the binding partner was pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the binding partner is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the binding partner (the antibody, in turn, may be directly labeled or indirectly labeled with a labeled anti-Ig antibody). Depending upon the order of addition of reaction components, test compounds which inhibit complex formation or which disrupt preformed complexes can be detected.

Alternatively, the reaction can be conducted in a liquid phase in the presence or absence of the test compound, the reaction products separated from unreacted components, and complexes detected; e.g., using an immobilized antibody specific for one binding partner to anchor any complexes formed in solution, and a labeled antibody specific for the other binding partner to detect anchored complexes. Again, depending upon the order of addition of reactants to the liquid phase, test compounds which inhibit complex or which disrupt preformed complexes can be identified.

In an alternate embodiment of the invention, a homogeneous assay can be used. In this approach, a preformed complex of the target gene protein and the interactive cellular or extracellular protein is prepared in which one of the binding partners is labeled, but the signal generated by the label is quenched due to complex formation (see, e.g., U.S. Patent No. 4,109,496 by Rubenstein which utilizes this approach for immunoassays). The addition of a test substance that competes with and displaces one of the binding partners from the preformed complex will result in the generation of a

signal above background. In this way, test substances which disrupt target gene protein-cellular or extracellular protein interaction can be identified.

In a particular embodiment, the target gene protein
5 can be prepared for immobilization using recombinant DNA techniques described in Section 5.4.2, *supra*. For example, the target gene coding region can be fused to a glutathione-S-transferase (GST) gene, using a fusion vector such as pGEX-5X-1, in such a manner that its binding activity is
10 maintained in the resulting fusion protein. The interactive cellular or extracellular protein can be purified and used to raise a monoclonal antibody, using methods routinely practiced in the art and described above, in Section 5.4.3. This antibody can be labeled with the radioactive isotope
15 ^{125}I , for example, by methods routinely practiced in the art. In a heterogeneous assay, e.g., the GST-target gene fusion protein can be anchored to glutathione-agarose beads. The interactive cellular or extracellular binding partner protein can then be added in the presence or absence of the test
20 compound in a manner that allows interaction and binding to occur. At the end of the reaction period, unbound material can be washed away, and the labeled monoclonal antibody can be added to the system and allowed to bind to the complexed binding partners. The interaction between the target gene
25 protein and the interactive cellular or extracellular binding partner protein can be detected by measuring the amount of radioactivity that remains associated with the glutathione-agarose beads. A successful inhibition of the interaction by the test compound will result in a decrease in measured
30 radioactivity.

Alternatively, the GST-target gene fusion protein and the interactive cellular or extracellular binding partner protein can be mixed together in liquid in the absence of the solid glutathione-agarose beads. The test compound can be
35 added either during or after the binding partners are allowed to interact. This mixture can then be added to the glutathione-agarose beads and unbound material is washed

fusion protein and allowed to bind. After washing away unbound peptides, labeled bound material, representing the cellular or extracellular binding partner protein binding domain, can be eluted, purified, and analyzed for amino acid
5 sequence by techniques well known in the art; e.g., using the Edman degradation procedure (see e.g., Creighton, 1983, Proteins: Structures and Molecular Principles, W.H. Freeman & Co., N.Y., pp. 34-49). Peptides so identified can be produced, using techniques well known in the art, either
10 synthetically (see e.g., Creighton, 1983, *supra* at pp. 50-60) or, if the gene has already been isolated, by using recombinant DNA technology, as described in Section 5.4.2, *supra*.

A particular embodiment of the invention features a
15 method of screening candidate compounds for their ability to antagonize the interaction between ligand and the receptor domain of a target gene product, including but not limited to the receptor domain of the rchd523 gene product. The rchd523 gene product, which is a G protein-coupled receptor protein
20 containing multiple transmembrane domains, is especially useful in screening for antagonists of ligand-receptor interactions. The method involves: a) mixing a candidate antagonist compound with a first compound which includes a recombinant target gene product comprising a receptor domain
25 (or ligand-binding fragment or analog) on the one hand and with a second compound which includes ligand on the other hand; b) determining whether the first and second compounds bind; and c) identifying antagonistic compounds as those which interfere with the binding of the first compound to the
30 second compound and/or which reduce the ligand-mediated release of intracellular Ca^{++} .

By an "antagonist" is meant a molecule which inhibits a particular activity, in this case, the ability of ligand to interact with a target gene product receptor domain
35 and/or to trigger the biological events resulting from such an interaction (e.g., release of intracellular Ca^{++}). Preferred therapeutics include antagonists, e.g., peptide

fragments (particularly, fragments derived from the N-terminal extracellular domain), antibodies (particularly, antibodies which recognize and bind the N-terminal extracellular domain), or drugs, which block ligand or target gene product function by interfering with the ligand-receptor interaction.

Because the receptor component of the target gene product can be produced by recombinant techniques and because candidate antagonists may be screened in vitro, the instant invention provides a simple and rapid approach to the identification of useful therapeutics.

Specific receptor fragments of interest include any portions of the target gene products that are capable of interaction with ligand, for example, all or part of the N-terminal extracellular domain. Such portions include the transmembrane segments and portions of the receptor deduced to be extracellular. Such fragments may be useful as antagonists (as described above), and are also useful as immunogens for producing antibodies which neutralize the activity of the target gene product in vivo (e.g., by interfering with the interaction between the receptor and ligand; see below). Extracellular regions may be identified by comparison with related proteins of similar structure (e.g., other members of the G-protein-coupled receptor superfamily); useful regions are those exhibiting homology to the extracellular domains of well-characterized members of the family.

Alternatively, from the primary amino acid sequence, the secondary protein structure and, therefore, the extracellular domain regions may be deduced semi-empirically using a hydrophobicity/hydrophilicity calculation such as the Chou-Fasman method (see, e.g., Chou and Fasman, *Ann. Rev. Biochem.* 47:251, 1978). Hydrophilic domains, particularly ones surrounded by hydrophobic stretches (e.g., transmembrane domains) present themselves as strong candidates for extracellular domains. Finally, extracellular domains may be

identified experimentally using standard enzymatic digest analysis, e.g., tryptic digest analysis.

Candidate fragments (e.g., all or part of the transmembrane segments or any extracellular fragment) are tested for interaction with ligand by the assays described herein (e.g., the assay described above). Such fragments are also tested for their ability to antagonize the interaction between ligand and its endogenous receptor using the assays described herein. Analogs of useful receptor fragments (as described above) may also be produced and tested for efficacy as screening components or antagonists (using the assays described herein); such analogs are also considered to be useful in the invention.

Of particular interest are receptor fragments encompassing the extracellular main-terminal domain (or a ligand binding fragment thereof). Also of interest are the target gene product extracellular loops. Peptide fragments derived from these extracellular loops may also be used as antagonists, particularly if the loops cooperate with the amino-terminal domain to facilitate ligand binding. Alternatively, such loops and extracellular N-terminal domain (as well as the full length target gene product) provide immunogens for producing anti-target gene product antibodies.

Binding of ligand to its receptor may be assayed by any of the methods described above in Section 5.5.1. Preferably, cells expressing recombinant target gene product (or a suitable target gene product fragment or analog) are immobilized on a solid substrate (e.g., the wall of a microtitre plate or a column) and reacted with detectably-labelled ligand (as described above). Binding is assayed by the detection label in association with the receptor component (and, therefore, in association with the solid substrate). Binding of labelled ligand to receptor-bearing cells is used as a "control" against which antagonist assays are measured. The antagonist assays involve incubation of the target gene product-bearing cells with an appropriate amount of candidate antagonist. To this mix, an equivalent

amount to labelled ligand is added. An antagonist useful in the invention specifically interferes with labelled ligand binding to the immobilized receptor-expressing cells.

An antagonist is then tested for its ability to
5 interfere with ligand function, i.e., to specifically interfere with labelled ligand binding without resulting in signal transduction normally mediated by the receptor. To test this using a functional assay, stably transfected cell lines containing the target gene product can be produced as
10 described herein and reporter compounds such as the calcium binding agent, FURA-2, loaded into the cytoplasm by standard techniques. Stimulation of the heterologous target gene product with ligand or another agonist leads to intracellular calcium release and the concomitant fluorescence of the
15 calcium-FURA-2 complex. This provides a convenient means for measuring agonist activity. Inclusion of potential antagonists along with ligand allows for the screening and identification of authentic receptor antagonists as those which effectively block ligand binding without producing
20 fluorescence (i.e., without causing the mobilization of intracellular Ca^{++}). Such an antagonist may be expected to be a useful therapeutic agent for cardiovascular disorders.

Appropriate candidate antagonists include target gene product fragments, particularly fragments containing a
25 ligand-binding portion adjacent to or including one or more transmembrane segments or an extracellular domain of the receptor (described above); such fragments would preferably include five or more amino acids. Other candidate antagonists include analogs of ligand and other peptides as
30 well as non-peptide compounds and anti-target gene product antibodies designed or derived from analysis of the receptor.

This screening method is described in detail with respect to the rchd523 gene in the example in Section 12, below. Because the rchd523 gene product is a G protein-
35 coupled receptor, antagonists of the interaction between the rchd523 gene product and its natural ligand provide excellent

candidates for compounds effective in the treatment of cardiovascular disease.

5.5.4. ASSAYS FOR AMELIORATION OF CARDIOVASCULAR DISEASE SYMPTOMS

5

Any of the binding compounds, including but not limited to compounds such as those identified in the foregoing assay systems, may be tested for the ability to ameliorate cardiovascular disease symptoms. Cell-based and animal model-based assays for the identification of compounds exhibiting such an ability to ameliorate cardiovascular disease symptoms are described below.

First, cell-based systems such as those described, above, in Section 5.4.4.2., may be used to identify compounds which may act to ameliorate cardiovascular disease symptoms. For example, such cell systems may be exposed to a compound, suspected of exhibiting an ability to ameliorate cardiovascular disease symptoms, at a sufficient concentration and for a time sufficient to elicit such an amelioration of cardiovascular disease symptoms in the exposed cells. After exposure, the cells are examined to determine whether one or more of the cardiovascular disease cellular phenotypes has been altered to resemble a more normal or more wild type, non-cardiovascular disease phenotype. For example, and not by way of limitation, in the case of monocytes, such more normal phenotypes may include but are not limited to decreased rates of LDL uptake, adhesion to endothelial cells, transmigration, foam cell formation, fatty streak formation, and production by foam cells of growth factors such as bFGF, IGF-I, VEGF, IL-1, M-CSF, TGF β , TGF α , TNF α , HB-EGF, PDGF, IFN- γ , and GM-CSF. Transmigration rates, for example, may be measured using the in vitro system of Navab et al., described in Section 5.1.1.3, above, by quantifying the number of monocytes that migrate across the endothelial monolayer and into the collagen layer of the subendothelial space.

Further, both cell-based systems and animal-based systems as described herein may be used to identify compounds which act to ameliorate symptoms of fibroproliferative and oncogenic related disorders, including tumorigenesis and the vascularization of tumors.

With regard to intervention, any treatments which reverse any aspect of symptoms of cardiovascular disease or fibroproliferative and oncogenic related disorders should be considered as candidates for human therapeutic intervention. Dosages of test agents may be determined by deriving dose-response curves, as discussed in Section 5.7.1, below.

Additionally, gene expression patterns may be
30 utilized to assess the ability of a compound to ameliorate
symptoms of cardiovascular disease or fibroproliferative and
oncogenic related disorders. For example, the expression
pattern of one or more fingerprint genes may form part of a
"fingerprint profile" which may be then be used in such an
35 assessment. "Fingerprint profile", as used herein, refers to
the pattern of mRNA expression obtained for a given tissue or
cell type under a given set of conditions. Such conditions

For example, and not by way of limitation, Paradigm A provides for the identification of fingerprint genes that are up-regulated in monocytes treated with oxidized LDL. Thus, to study the effect of anti-oxidant drugs, for example, 5 in a clinical trial, blood may be drawn from patients before and at different stages during treatment with such a drug. Their monocytes may then be isolated and RNA prepared and analyzed by differential display as described in Sections 6.1.1 and 6.1.2. The levels of expression of these 10 fingerprint genes may be quantified by Northern blot analysis or RT-PCR, as described in Section 6.1.2, or by one of the methods described in Section 5.8.1, or alternatively by measuring the amount of protein produced, by one of the methods described in Section 5.8.2. In this way, the 15 fingerprint profiles may serve as surrogate markers indicative of the physiological response of monocytes that have taken up oxidized LDL. Accordingly, this response state may be determined before, and at various points during, drug treatment. This method is described in further detail in the 20 example in Section 10, below.

This method may also be applied to the other paradigms disclosed herein. For example, and not by way of limitation, the fingerprint profile of Paradigm B reveals that bcl-2 and glutathione peroxidase are both down-regulated 25 in the monocytes of patients exposed to a high lipid diet, e.g. cholesterol or fat, that leads to high serum LDL levels. Drugs may be tested, for example, for their ability to ameliorate the effects of hypercholesterolemia in clinical trials. Patients with high LDL levels may have their 30 monocytes isolated before, and at different stages after, drug treatment. The drug's efficacy may be measured by determining the degree of restored expression of bcl-2 and glutathione peroxidase, as described above for the Paradigm A fingerprint profile.

35

5.5.6. ASSAYS FOR COMPOUNDS THAT MODULATE
EXPRESSION OF TARGET GENES

Compounds and other substances that modulate expression of target genes can be screened using in vitro cellular systems. In a manner analogous to the monitoring of compounds clinical samples described in Section 5.5.5, above, a sample of cells, such as a tissue culture is exposed to a test substance. Appropriate tissue culture cells include, but are not limited to, human umbilical vein endothelial cells (HUVECs), bovine aortic endothelial cells (BAECs), and 293 cells (embryonic human kidney cells). The RNA is then extracted from the cells. The level of transcription of a specific target gene can be detected using, for example, standard RT-PCR amplification techniques and/or Northern analysis (as described in the example in Section 6.1.2, below). Alternatively, the level of target protein production can be assayed by using antibodies that detect the target gene protein, as described in Section 5.8.2, below. The level of expression is compared to a control cell sample which was not exposed to the test substance.

Compounds that can be screened for modulation of expression of the target gene include, but are not limited to, small inorganic or organic molecules, peptides, such as peptide hormones analogs, steroid hormones, analogs of such hormones, and other proteins. Compounds that down-regulate expression include, but are not limited to, oligonucleotides that are complementary to the 5'-end of the mRNA of the target gene and inhibit transcription by forming triple helix structures, and ribozymes or antisense molecules which inhibit translation of the target gene mRNA. Techniques and strategies for designing such down-regulating test compounds are described in detail in Section 5.6, below.

5.6. COMPOUNDS AND METHODS FOR TREATMENT OF
CARDIOVASCULAR DISEASE

Described below are methods and compositions whereby cardiovascular disease symptoms may be ameliorated. The methods and compositions described below may also be

applied to the amelioration of symptoms associated with fibroproliferative and oncogenic disorders, but by way of example, and not by limitation, are described in the subsections below in terms of cardiovascular disease.

- 5 Certain cardiovascular diseases are brought about, at least in part, by an excessive level of gene product, or by the presence of a gene product exhibiting an abnormal or excessive activity. As such, the reduction in the level and/or activity of such gene products would bring about the
- 10 amelioration of cardiovascular disease symptoms. Techniques for the reduction of target gene expression levels or target gene product activity levels are discussed in Section 5.6.1, below.

- Alternatively, certain other cardiovascular
- 15 diseases are brought about, at least in part, by the absence or reduction of the level of gene expression, or a reduction in the level of a gene product's activity. As such, an increase in the level of gene expression and/or the activity of such gene products would bring about the amelioration of
- 20 cardiovascular disease symptoms.

- In some cases, the up-regulation of a gene in a disease state reflects a protective role for that gene product in responding to the disease condition. Enhancement of such a target gene's expression, or the activity of the
- 25 target gene product, will reinforce the protective effect it exerts. Some cardiovascular disease states may result from an abnormally low level of activity of such a protective gene. In these cases also, an increase in the level of gene expression and/or the activity of such gene products would
- 30 bring about the amelioration of cardiovascular disease symptoms. Techniques for increasing target gene expression levels or target gene product activity levels are discussed in Section 5.6.2, below.

5.6.1. COMPOUNDS THAT INHIBIT EXPRESSION,
SYNTHESIS OR ACTIVITY OF MUTANT TARGET
GENE ACTIVITY

As discussed above, target genes involved in cardiovascular disease disorders can cause such disorders via an increased level of target gene activity. As summarized in Table 1, above, and detailed in the examples in Sections 8 and 9, below, a number of genes are now known to be up-regulated in endothelial cells under disease conditions. Specifically, rchd005, rchd024, rchd032, and rchd036 are all up-regulated in endothelial cells treated with IL-1. Furthermore, rchd502, rchd523, rchd528, rchd534, COX II, and MnSOD are all up-regulated in endothelial cells subjected to shear stress. In some cases, such up-regulation may have a causative or exacerbating effect on the disease state. A variety of techniques may be utilized to inhibit the expression, synthesis, or activity of such target genes and/or proteins.

For example, compounds such as those identified through assays described, above, in Section 5.5, which exhibit inhibitory activity, may be used in accordance with the invention to ameliorate cardiovascular disease symptoms. As discussed in Section 5.5, above, such molecules may include, but are not limited to small organic molecules, peptides, antibodies, and the like. Inhibitory antibody techniques are described, below, in Section 5.6.1.2.

For example, compounds can be administered that compete with endogenous ligand for the rchd523 gene product. The resulting reduction in the amount of ligand-bound rchd523 gene transmembrane protein will modulated endothelial cell physiology. Compounds that can be particularly useful for this purpose include, for example, soluble proteins or peptides, such as peptides comprising one or more of the extracellular domains, or portions and/or analogs thereof, of the rchd523 gene product, including, for example, soluble fusion proteins such as Ig-tailed fusion proteins. (For a discussion of the production of Ig-tailed fusion proteins, see, for example, U.S. Patent No. 5,116,964.).

Alternatively, compounds, such as ligand analogs or antibodies, that bind to the rchd523 gene product receptor site, but do not activate the protein, (e.g., receptor-ligand antagonists) can be effective in inhibiting rchd523 gene product activity.

Further, antisense and ribozyme molecules which inhibit expression of the target gene may also be used in accordance with the invention to inhibit the aberrant target gene activity. Such techniques are described, below, in Section 5.6.1.1. Still further, also as described, below, in Section 5.6.1.1, triple helix molecules may be utilized in inhibiting the aberrant target gene activity.

5.6.1.1. INHIBITORY ANTISENSE, RIBOZYME, TRIPLE HELIX, AND GENE INACTIVATION APPROACHES

Among the compounds which may exhibit the ability to ameliorate cardiovascular disease symptoms are antisense, ribozyme, and triple helix molecules. Such molecules may be designed to reduce or inhibit mutant target gene activity. Techniques for the production and use of such molecules are well known to those of skill in the art.

Antisense RNA and DNA molecules act to directly block the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation.

Antisense approaches involve the design of oligonucleotides (either DNA or RNA) that are complementary to target gene mRNA. The antisense oligonucleotides will bind to the complementary target gene mRNA transcripts and prevent translation. Absolute complementarity, although preferred, is not required. A sequence "complementary" to a portion of an RNA, as referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double-stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense

nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with an RNA it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

Oligonucleotides that are complementary to the 5' end of the message, e.g., the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have recently shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., 1994, Nature 372:333-335. Thus, oligonucleotides complementary to either the 5'- or 3'- non-translated, non-coding regions of the target gene could be used in an antisense approach to inhibit translation of endogenous target gene mRNA. Oligonucleotides complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon.

Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the 5'-, 3'- or coding region of target gene mRNA, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

30 Regardless of the choice of target sequence, it is preferred that *in vitro* studies are first performed to quantitate the ability of the antisense oligonucleotide to inhibit gene expression. It is preferred that these studies utilize controls that distinguish between antisense gene
35 inhibition and nonspecific biological effects of oligonucleotides. It is also preferred that these studies compare levels of the target RNA or protein with that of an

internal control RNA or protein. Additionally, it is envisioned that results obtained using the antisense oligonucleotide are compared with those obtained using a control oligonucleotide. It is preferred that the control
5 oligonucleotide is of approximately the same length as the test oligonucleotide and that the nucleotide sequence of the oligonucleotide differs from the antisense sequence no more than is necessary to prevent specific hybridization to the target sequence.

10 The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, 15 hybridization, etc. The oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors in vivo), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al., 1989, Proc. Natl. Acad. Sci. U.S.A. 86:6553-6556; Lemaitre et al., 20 1987, Proc. Natl. Acad. Sci. 84:648-652; PCT Publication No. WO88/09810, published December 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication No. WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents. (See, e.g., Krol et al., 1988, BioTechniques 6:958-976) or 25 intercalating agents. (See, e.g., Zon, 1988, Pharm. Res. 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

1-methylguanine, 1-methylinosine, 2,2-dimethylguanine,
 2-methyladenine, 2-methylguanine, 3-methylcytosine,
 5-methylcytosine, N6-adenine, 7-methylguanine,
 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil,
 5 beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil,
 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine,
 uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil,
 queosine, 2-thiocytosine, 5-methyl-2-thiouracil,
 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-
 10 5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v),
 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl)
 uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide may also comprise at
 least one modified sugar moiety selected from the group
 15 including but not limited to arabinose, 2-fluoroarabinose,
 xylulose, and hexose.

In yet another embodiment, the antisense
 oligonucleotide comprises at least one modified phosphate
 backbone selected from the group consisting of a
 20 phosphorothioate, a phosphorodithioate, a
 phosphoramidothioate, a phosphoramidate, a phosphordiamidate,
 a methylphosphonate, an alkyl phosphotriester, and a
 formacetal or analog thereof.

In yet another embodiment, the antisense
 25 oligonucleotide is an α -anomeric oligonucleotide. An α -
 anomeric oligonucleotide forms specific double-stranded
 hybrids with complementary RNA in which, contrary to the
 usual β -units, the strands run parallel to each other
 (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641). The
 30 oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al.,
 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA
 analogue (Inoue et al., 1987, FEBS Lett. 215:327-330).

Oligonucleotides of the invention may be
 synthesized by standard methods known in the art, e.g. by use
 35 of an automated DNA synthesizer (such as are commercially
 available from Biosearch, Applied Biosystems, etc.). As
 examples, phosphorothioate oligonucleotides may be

synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-5 7451), etc.

While antisense nucleotides complementary to the target gene coding region sequence could be used, those complementary to the transcribed untranslated region are most preferred.

10 Specific antisense oligonucleotides for the rchd534 gene (both short and long spliceoforms) and fchd540 gene are described in the example in Section 17, below.

 The antisense molecules should be delivered to cells which express the target gene *in vivo*, e.g.,
15 endothelial cells. A number of methods have been developed for delivering antisense DNA or RNA to cells; e.g., antisense molecules can be injected directly into the tissue site, or modified antisense molecules, designed to target the desired cells (e.g., antisense linked to peptides or antibodies that
20 specifically bind receptors or antigens expressed on the target cell surface) can be administered systemically.

 However, it is often difficult to achieve intracellular concentrations of the antisense sufficient to suppress translation of endogenous mRNAs. Therefore a
25 preferred approach utilizes a recombinant DNA construct in which the antisense oligonucleotide is placed under the control of a strong pol III or pol II promoter. The use of such a construct to transfect target cells in the patient will result in the transcription of sufficient amounts of
30 single stranded RNAs that will form complementary base pairs with the endogenous target gene transcripts and thereby prevent translation of the target gene mRNA. For example, a vector can be introduced in vivo such that it is taken up by a cell and directs the transcription of an antisense RNA.
35 Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by

recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in mammalian cells. Expression of the sequence encoding the antisense RNA can be
5 by any promoter known in the art to act in mammalian, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include but are not limited to: the SV40 early promoter region (Bernoist and Chambon, 1981, Nature 290:304-310), the promoter contained in the 3' long
10 terminal repeat of Rous sarcoma virus (Yamamoto et al., 1980, Cell 22:787-797), the herpes thymidine kinase promoter (Wagner et al., 1981, Proc. Natl. Acad. Sci. U.S.A. 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster et al., 1982, Nature 296:39-42), etc. Any type of
15 plasmid, cosmid, YAC or viral vector can be used to prepare the recombinant DNA construct which can be introduced directly into the tissue site; e.g., atherosclerotic vascular tissue. Alternatively, viral vectors can be used which selectively infect the desired tissue, in which case
20 administration may be accomplished by another route (e.g., systemically).

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. The mechanism of ribozyme action involves sequence specific hybridization of
25 the ribozyme molecule to complementary target RNA, followed by an endonucleolytic cleavage. Ribozyme molecules designed to catalytically cleave target gene mRNA transcripts can also be used to prevent translation of target gene mRNA and expression of target gene. (See, e.g., PCT International
30 Publication WO90/11364, published October 4, 1990; Sarver et al., 1990, Science 247:1222-1225). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy target gene mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at
35 locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following

sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, 1988, Nature, 334:585-591. For example, there are hundreds of
5 potential hammerhead ribozyme cleavage sites within the nucleotide sequence of rchd534 (both short and long spliceoforms) and fchd540 cDNA. Preferably the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the target mRNA; i.e., to increase
10 efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

Specific hammerhead ribozymes molecules for the rchd534 (both short and long spliceoforms) and fchd540 genes are described in the Example in Section 13, below.

15 The ribozymes of the present invention also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as the one which occurs naturally in Tetrahymena Thermophila (known as the IVS, or L-19 IVS RNA) and which has been extensively described by Thomas Cech and collaborators
20 (Zaug, et al., 1984, Science, 224:574-578; Zaug and Cech, 1986, Science, 231:470-475; Zaug, et al., 1986, Nature, 324:429-433; published International patent application No. WO 88/04300 by University Patents Inc.; Been and Cech, 1986, Cell, 47:207-216). The Cech-type ribozymes have an eight
25 base pair active site which hybridizes to a target RNA sequence whereafter cleavage of the target RNA takes place. The invention encompasses those Cech-type ribozymes which target eight base-pair active site sequences that are present in target gene.

30 As in the antisense approach, the ribozymes can be composed of modified oligonucleotides (e.g. for improved stability, targeting, etc.) and should be delivered to cells which express the target gene *in vivo*, e.g., endothelial cells. A preferred method of delivery involves using a DNA
35 construct "encoding" the ribozyme under the control of a strong constitutive pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the

maintained, nucleic acid molecules that encode and express target gene polypeptides exhibiting normal activity may be introduced into cells via gene therapy methods such as those described, below, in Section 5.7. that do not contain

- 5 sequences susceptible to whatever antisense, ribozyme, or triple helix treatments are being utilized. Alternatively, it may be preferable to coadminister normal target gene protein into the cell or tissue in order to maintain the requisite level of cellular or tissue target gene activity.
- 10 Endogenous target gene expression can also be reduced by inactivating or "knocking out" the target gene or its promoter using targeted homologous recombination. (E.g., see Smithies et al., 1985, Nature 317:230-234; Thomas & Capecchi, 1987, Cell 51:503-512; Thompson et al., 1989 Cell
- 15 5:313-321; each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional target (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous target gene (either the coding regions or regulatory regions of the target gene) can
- 20 be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express target *in vivo*. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the target gene. Such approaches can be adapted for use in
- 25 humans provided the recombinant DNA constructs are directly administered or targeted to the required site in vivo using appropriate viral vectors, e.g., vectors for delivery vascular tissue.

- Alternatively, endogenous target gene expression
- 30 can be reduced by targeting deoxyribonucleotide sequences complementary to the regulatory region of the target gene (i.e., the target promoter and/or enhancers) to form triple helical structures that prevent transcription of the target gene in target cells in the body. (See generally, Helene, C.
- 35 1991, Anticancer Drug Des., 6(6):569-84; Helene, C., et al., 1992, Ann, N.Y. Accad. Sci., 660:27-36; and Maher, L.J., 1992, Bioassays 14(12):807-15).

intracellular target gene epitopes may also be administered. Such single chain antibodies may be administered, for example, by expressing nucleotide sequences encoding single-chain antibodies within the target cell population by
5 utilizing, for example, techniques such as those described in Marasco et al. (Marasco, W. et al., 1993, Proc. Natl. Acad. Sci. USA 90:7889-7893).

In some instances, the target gene protein is extracellular, or is a transmembrane protein, such as the rchd523 gene product. Antibodies that are specific for one or more extracellular domains of the rchd523 gene product, for example, and that interfere with its activity, are particularly useful in treating cardiovascular disease. Such antibodies are especially efficient because they can access the target domains directly from the bloodstream. Any of the administration techniques described, below in Section 5.7 which are appropriate for peptide administration may be utilized to effectively administer inhibitory target gene antibodies to their site of action.

5.6.2. METHODS FOR RESTORING OR ENHANCING TARGET GENE ACTIVITY

Target genes that cause cardiovascular disease may be underexpressed within cardiovascular disease situations.

25 As summarized in Table 1, above, and detailed in the example in Sections 7, below, several genes are now known to be down-regulated in monocytes under disease conditions.

Specifically, bcl-2 and glutathione peroxidase gene expression is down-regulated in the monocytes of patients

30 exposed to a high lipid diet, e.g. cholesterol or fat, that leads to high serum LDL levels. Alternatively, the activity of target gene products may be decreased, leading to the development of cardiovascular disease symptoms. Such down-regulation of target gene expression or decrease of target

35 gene product activity might have a causative or exacerbating effect on the disease state.

In some cases, target genes that are up-regulated in the disease state might be exerting a protective effect. As summarized in Table 1, above, and detailed in the examples in Sections 8 and 9, below, a number of genes are now known
5 to be up-regulated in endothelial cells under disease conditions. Specifically, rchd005, rchd024, rchd032, and rchd036 are all up-regulated in endothelial cells treated with IL-1. Furthermore, rchd502, rchd523, rchd528, rchd534, COX II, and MnSOD are all up-regulated in endothelial cells
10 subjected to shear stress. A variety of techniques may be utilized to increase the expression, synthesis, or activity of such target genes and/or proteins, for those genes that exert a protective effect in response to disease conditions.

Described in this Section are methods whereby the
15 level of target gene activity may be increased to levels wherein cardiovascular disease symptoms are ameliorated. The level of gene activity may be increased, for example, by either increasing the level of target gene product present or by increasing the level of active target gene product which
20 is present.

For example, a target gene protein, at a level sufficient to ameliorate cardiovascular disease symptoms may be administered to a patient exhibiting such symptoms. Any of the techniques discussed, below, in Section 5.7, may be
25 utilized for such administration. One of skill in the art will readily know how to determine the concentration of effective, non-toxic doses of the normal target gene protein, utilizing techniques such as those described, below, in Section 5.7.1.

30 Additionally, RNA sequences encoding target gene protein may be directly administered to a patient exhibiting cardiovascular disease symptoms, at a concentration sufficient to produce a level of target gene protein such that cardiovascular disease symptoms are ameliorated. Any of
35 the techniques discussed, below, in Section 5.7, which achieve intracellular administration of compounds, such as, for example, liposome administration, may be utilized for the

administration of such RNA molecules. The RNA molecules may be produced, for example, by recombinant techniques such as those described, above, in Section 5.4.2.

- Further, patients may be treated by gene replacement therapy. One or more copies of a normal target gene, or a portion of the gene that directs the production of a normal target gene protein with target gene function, may be inserted into cells using vectors which include, but are not limited to adenovirus, adeno-associated virus, and retrovirus vectors, in addition to other particles that introduce DNA into cells, such as liposomes. Additionally, techniques such as those described above may be utilized for the introduction of normal target gene sequences into human cells.
- Cells, preferably, autologous cells, containing normal target gene expressing gene sequences may then be introduced or reintroduced into the patient at positions which allow for the amelioration of cardiovascular disease symptoms. Such cell replacement techniques may be preferred, for example, when the target gene product is a secreted, extracellular gene product.

5.7. PHARMACEUTICAL PREPARATIONS AND METHODS OF ADMINISTRATION

- The identified compounds that inhibit target gene expression, synthesis and/or activity can be administered to a patient at therapeutically effective doses to treat or ameliorate cardiovascular disease. A therapeutically effective dose refers to that amount of the compound sufficient to result in amelioration of symptoms of cardiovascular disease.

5.7.1. EFFECTIVE DOSE

- Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the

ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Compounds which exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

10 The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may
15 vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal
20 models to achieve a circulating plasma concentration range that includes the IC₅₀ (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in
25 humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

5.7.2. FORMULATIONS AND USE

Pharmaceutical compositions for use in accordance
30 with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers or excipients.

Thus, the compounds and their physiologically acceptable salts and solvates may be formulated for
35 administration by inhalation or insufflation (either through the mouth or the nose) or oral, buccal, parenteral or rectal administration.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g.,
5 pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch
10 glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water
15 or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin
20 or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavoring, coloring and sweetening agents as
25 appropriate.

Preparations for oral administration may be suitably formulated to give controlled release of the active compound.

For buccal administration the compositions may take
30 the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from
35 pressurized packs or a nebuliser, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon

dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be
5 formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be
10 presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain
15 formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, e.g.,
20 containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be
25 administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as
30 sparingly soluble derivatives, for example, as a sparingly soluble salt.

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may
35 for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration.

5.8. DIAGNOSIS OF CARDIOVASCULAR DISEASE ABNORMALITIES

A variety of methods may be employed, utilizing reagents such as fingerprint gene nucleotide sequences described in Section 5.4.1, and antibodies directed against 5 differentially expressed and pathway gene peptides, as described, above, in Sections 5.4.2. (peptides) and 5.4.3. (antibodies). Specifically, such reagents may be used, for example, for the detection of the presence of target gene mutations, or the detection of either over or under 10 expression of target gene mRNA.

The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one specific fingerprint gene nucleic acid or anti-fingerprint gene antibody reagent described herein, which may 15 be conveniently used, e.g., in clinical settings, to diagnose patients exhibiting cardiovascular disease symptoms or at risk for developing cardiovascular disease.

Any cell type or tissue, preferably monocytes, endothelial cells, or smooth muscle cells, in which the 20 fingerprint gene is expressed may be utilized in the diagnostics described below.

5.8.1. DETECTION OF FINGERPRINT GENE NUCLEIC ACIDS

25 DNA or RNA from the cell type or tissue to be analyzed may easily be isolated using procedures which are well known to those in the art. Diagnostic procedures may also be performed "in situ" directly upon tissue sections (fixed and/or frozen) of patient tissue obtained from 30 biopsies or resections, such that no nucleic acid purification is necessary. Nucleic acid reagents such as those described in Section 5.1. may be used as probes and/or primers for such in situ procedures (see, for example, Nuovo, G.J., 1992, PCR in situ hybridization: protocols and 35 applications, Raven Press, NY).

Fingerprint gene nucleotide sequences, either RNA or DNA, may, for example, be used in hybridization or

amplification assays of biological samples to detect cardiovascular disease-related gene structures and expression. Such assays may include, but are not limited to, Southern or Northern analyses, single stranded conformational polymorphism analyses, in situ hybridization assays, and polymerase chain reaction analyses. Such analyses may reveal both quantitative aspects of the expression pattern of the fingerprint gene, and qualitative aspects of the fingerprint gene expression and/or gene composition. That is, such aspects may include, for example, point mutations, insertions, deletions, chromosomal rearrangements, and/or activation or inactivation of gene expression.

Such an in situ hybridization analysis is described in the example in Section 14, below. Specifically, high levels of expression of the rchd502 and rchd528 genes were detected specifically within the endothelial cells of diseased tissue removed from a human cardiovascular disease patient, and not in any other cell type present in the tissue, including smooth muscle cells and macrophages. These results clearly demonstrate how the target genes described herein provide for novel diagnoses of cardiovascular disease. Furthermore, because these diagnoses are correlated with specific target genes, they allow for more specifically directed methods of treatment of cardiovascular disease.

Preferred diagnostic methods for the detection of fingerprint gene-specific nucleic acid molecules may involve for example, contacting and incubating nucleic acids, derived from the cell type or tissue being analyzed, with one or more labeled nucleic acid reagents as are described in Section 5.1, under conditions favorable for the specific annealing of these reagents to their complementary sequences within the nucleic acid molecule of interest. Preferably, the lengths of these nucleic acid reagents are at least 9 to 30 nucleotides. After incubation, all non-annealed nucleic acids are removed from the nucleic acid:fingerprint molecule hybrid. The presence of nucleic acids from the fingerprint tissue which have hybridized, if any such molecules exist, is

primers) in the reverse transcription and nucleic acid amplification steps of this method are chosen from among the fingerprint gene nucleic acid reagents described in Section 5.1. The preferred lengths of such nucleic acid reagents are
5 at least 15-30 nucleotides. For detection of the amplified product, the nucleic acid amplification may be performed using radioactively or non-radioactively labeled nucleotides. Alternatively, enough amplified product may be made such that the product may be visualized by standard ethidium bromide
10 staining or by utilizing any other suitable nucleic acid staining method.

In addition to methods which focus primarily on the detection of one nucleic acid sequence, fingerprint profiles, as discussed in Section 5.5.4, may also be assessed in such
15 detection schemes. Fingerprint profiles may be generated, for example, by utilizing a differential display procedure, as discussed, above, in Section 5.1.2, Northern analysis and/or RT-PCR. Any of the gene sequences described, above, in Section 5.4.1. may be used as probes and/or PCR primers
20 for the generation and corroboration of such fingerprint profiles.

5.8.2. DETECTION OF FINGERPRINT GENE PEPTIDES

Antibodies directed against wild type or mutant
25 fingerprint gene peptides, which are discussed, above, in Section 5.4.3, may also be used as cardiovascular disease diagnostics and prognostics, as described, for example, herein. Such diagnostic methods, may be used to detect abnormalities in the level of fingerprint gene protein
30 expression, or abnormalities in the structure and/or tissue, cellular, or subcellular location of fingerprint gene protein. Structural differences may include, for example, differences in the size, electronegativity, or antigenicity of the mutant fingerprint gene protein relative to the normal
35 fingerprint gene protein.

Protein from the tissue or cell type to be analyzed may easily be detected or isolated using techniques which are

well known to those of skill in the art, including but not limited to western blot analysis. For a detailed explanation of methods for carrying out western blot analysis, see Sambrook et al, 1989, supra, at Chapter 18. The protein
5 detection and isolation methods employed herein may also be such as those described in Harlow and Lane, for example, (Harlow, E. and Lane, D., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York), which is incorporated herein by reference
10 in its entirety.

Preferred diagnostic methods for the detection of wild type or mutant fingerprint gene peptide molecules may involve, for example, immunoassays wherein fingerprint gene peptides are detected by their interaction with an anti-
15 fingerprint gene specific peptide antibody.

For example, antibodies, or fragments of antibodies, such as those described, above, in Section 5.4.3, useful in the present invention may be used to quantitatively or qualitatively detect the presence of wild type or mutant
20 fingerprint gene peptides. This can be accomplished, for example, by immunofluorescence techniques employing a fluorescently labeled antibody (see below) coupled with light microscopic, flow cytometric, or fluorimetric detection. Such techniques are especially preferred if the fingerprint
25 gene peptides are expressed on the cell surface.

The antibodies (or fragments thereof) useful in the present invention may, additionally, be employed histologically, as in immunofluorescence or immunoelectron microscopy, for in situ detection of fingerprint gene
30 peptides. In situ detection may be accomplished by removing a histological specimen from a patient, and applying thereto a labeled antibody of the present invention. The antibody (or fragment) is preferably applied by overlaying the labeled antibody (or fragment) onto a biological sample. Through the
35 use of such a procedure, it is possible to determine not only the presence of the fingerprint gene peptides, but also their distribution in the examined tissue. Using the present

invention, those of ordinary skill will readily perceive that any of a wide variety of histological methods (such as staining procedures) can be modified in order to achieve such in situ detection.

- 5 Immunoassays for wild type or mutant fingerprint gene peptides typically comprise incubating a biological sample, such as a biological fluid, a tissue extract, freshly harvested cells, or cells which have been incubated in tissue culture, in the presence of a detectably labeled antibody
- 10 capable of identifying fingerprint gene peptides, and detecting the bound antibody by any of a number of techniques well known in the art.

- The biological sample may be brought in contact with and immobilized onto a solid phase support or carrier
- 15 such as nitrocellulose, or other solid support which is capable of immobilizing cells, cell particles or soluble proteins. The support may then be washed with suitable buffers followed by treatment with the detectably labeled fingerprint gene specific antibody. The solid phase support
- 20 may then be washed with the buffer a second time to remove unbound antibody. The amount of bound label on solid support may then be detected by conventional means.

- By "solid phase support or carrier" is intended any support capable of binding an antigen or an antibody. Well-
- 25 known supports or carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble for the purposes of the
- 30 present invention. The support material may have virtually any possible structural configuration so long as the coupled molecule is capable of binding to an antigen or antibody. Thus, the support configuration may be spherical, as in a bead, or cylindrical, as in the inside surface of a test
- 35 tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Preferred supports include polystyrene beads. Those skilled

3640E0" 9824E050

Detection may also be accomplished using any of a variety of other immunoassays. For example, by radioactively labeling the antibodies or antibody fragments, it is possible to detect fingerprint gene wild type or mutant peptides through the use of a radioimmunoassay (RIA) (see, for example, Weintraub, B., *Principles of Radioimmunoassays*, Seventh Training Course on Radioligand Assay Techniques, The Endocrine Society, March, 1986, which is incorporated by reference herein). The radioactive isotope can be detected by such means as the use of a gamma counter, or a scintillation counter or by autoradiography.

It is also possible to label the antibody with a fluorescent compound. When the fluorescently labeled antibody is exposed to light of the proper wave length, its presence can then be detected due to fluorescence. Among the most commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine.

The antibody can also be detectably labeled using fluorescence emitting metals such as ¹⁵²Eu, or others of the lanthanide series. These metals can be attached to the antibody using such metal chelating groups as diethylenetriaminepentacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA).

The antibody also can be detectably labeled by coupling it to a chemiluminescent compound. The presence of the chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of particularly useful chemiluminescent labeling compounds are luminol, isoluminol, theromatic acridinium ester, imidazole, acridinium salt and oxalate ester.

Likewise, a bioluminescent compound may be used to label the antibody of the present invention. Bioluminescence is a type of chemiluminescence found in biological systems in, which a catalytic protein increases the efficiency of the

imaging techniques. An antibody specific for a target gene product, or preferably an antigen binding fragment thereof, is conjugated to a label (e.g., a gamma emitting radioisotope) which generates a detectable signal and
5 administered to a subject (human or animal) suspected of having cardiovascular disease. After sufficient time to allow the detectably-labeled antibody to localize at the diseased or damaged tissue site (or sites), the signal generated by the label is detected by a photoscanning device.
10 The detected signal is then converted to an image of the tissue. This image makes it possible to localize the tissue in vivo. This data can then be used to develop an appropriate therapeutic strategy.

Antibody fragments, rather than whole antibody
15 molecules, are generally preferred for use in tissue imaging. Antibody fragments accumulate at the tissue(s) more rapidly because they are distributed more readily than are entire antibody molecules. Thus an image can be obtained in less time than is possible using whole antibody. These fragments
20 are also cleared more rapidly from tissues, resulting in a lower background signal. See, e.g., Haber et al., U.S. Patent No. 4,036,945; Goldenberg et al., U.S. Patent No. 4,331,647. The divalent antigen binding fragment (Fab')₂ and the monovalent Fab are especially preferred. Such fragments
25 can be prepared by digestion of the whole immunoglobulin molecule with the enzymes pepsin or papain according to any of several well known protocols. The types of labels that are suitable for conjugation to a monoclonal antibody for diseased or damaged tissue localization include, but are not
30 limited to radiolabels (i.e., radioisotopes), fluorescent labels and biotin labels.

Among the radioisotopes that can be used to label antibodies or antibody fragments, gamma-emitters, positron-emitters, X-ray-emitters and fluorescence-emitters are
35 suitable for localization. Suitable radioisotopes for labeling antibodies include Iodine-131, Iodine-123, Iodine-125, Iodine-126, Iodine-133, Bromine-77, Indium-111, Indium-

113m, Gallium-67, Gallium-68, Ruthenium-95, Ruthenium-97, Ruthenium-103, Ruthenium-105, Mercury-107, Mercury-203, Rhenium-99m, Rhenium-105, Rhenium-101, Tellurium-121m, Tellurium-122m, Tellurium-125m, Thulium-165, Thulium-167,
5 Thulium-168, Technetium-99m and Fluorine-18. The halogens can be used more or less interchangeably as labels since halogen-labeled antibodies and/or normal immunoglobulins would have substantially the same kinetics and distribution and similar metabolism.

10 The gamma-emitters Indium-111 and Technetium-99m are preferred because these radiometals are detectable with a gamma camera and have favorable half lives for imaging in vivo. Antibody can be labelled with Indium-111 or Technetium-99m via a conjugated metal chelator, such as DTPA
15 (diethylenetriaminepentaacetic acid). See Krejcarek et al., 1977, Biochem. Biophys. Res. Comm. 77:581; Khaw et al., 1980, Science 209:295; Gansow et al., U.S. Patent No. 4,472,509; Hnatowich, U.S. Patent No. 4,479,930, the teachings of which are incorporated herein by reference.

20 Fluorescent compounds that are suitable for conjugation to a monoclonal antibody include fluorescein sodium, fluorescein isothiocyanate, and Texas Red sulfonyl chloride. See, DeBelder & Wik, 1975, Carbohydrate Research 44:254-257. Those skilled in the art will know, or will be
25 able to ascertain with no more than routine experimentation, other fluorescent compounds that are suitable for labeling monoclonal antibodies.

6. EXAMPLE: IDENTIFICATION OF GENES DIFFERENTIALLY
30 EXPRESSED IN RESPONSE TO PARADIGM A: IN VITRO FOAM CELL
 PARADIGM

 According to the invention, differential display may be used to detect genes that are differentially expressed in monocytes that were treated so as to simulate the
35 conditions under which foam cells develop during atherogenesis.

6.1. MATERIALS AND METHODS

6.1.1. CELL ISOLATION AND CULTURING

Blood (~200 ml) was drawn into chilled 20 ml vacutainer tubes to which 3 ml of citrate phosphate dextrose (Sigma) was added. Blood was then pooled into 50 ml tubes and spun in the Beckman GS-6R at 1250 RPM for 15 minutes at 4°C. The upper clear layer (~25 ml) was then removed with a pipette and discarded and replaced with the same volume of 4°C PBS. The blood was then mixed, and spun again at 2680 RPM for 15 minutes at 4°C. The upper layer was then removed and discarded, and the buffy coat at the interface was removed in ~5 ml and placed in a separate 50 ml tube, and the pipette was washed with 20 ml PBS. Cells were added to a T flask and stored at 4°C for 16 hours. A small aliquot of the cells were then removed and counted using a hemacytometer. The final red blood cell concentration in the buffy coat population was then adjusted to 1.5×10^9 /ml with PBS, the cells were added to Leucoprep tubes (Becton Dickinson) after being allowed to come to room temperature, and spun at 2300 RPM for 25 minutes at 25°C. The upper clear layer was removed and discarded and the turbid layer over the gel was removed and pooled in 50 ml tubes. Samples were then diluted to 50 ml with PBS (25°C) and spun at 1000 RPM for 10 minutes. The supernatant was then removed, and the pellet was resuspended in 50 ml PBS. This procedure was repeated 3 more times. After the last spin, the cells were resuspended in a small volume of PBS and counted.

Tissue culture dishes were coated with bovine collagen before monocytes were plated out. 1/6 volume of 7X RPMI (JRH Biosciences) was added to Vitrogen 100 collagen (Celtrix) which was then diluted 1:10 with RPMI to a final concentration of 0.35 mg/ml. Collagen mixture was then added to plates (2.5 ml/100 mm dish) and placed at 37°C for at least one hour to allow for gel formation. After gel formation has taken place, the RPMI was removed and cells were added in RPMI/10% plasma derived serum (PDS). PDS was prepared by drawing blood into chilled evacuated tubes

containing 1/10th volume 3.8% sodium citrate. Blood was then transferred into new Sorvall tubes and spun at 14,000-16,000 RPM for 20 minutes at 4°C. Plasma layer was removed and pooled in new tubes to which 1/50th volume 1M CaCl₂ was added. Plasma was mixed and aliquoted into new Sorvall tubes and incubated at 37° for 2 hours to allow for fibrin clot formation. The clot was then disturbed with a pipette to allow it to contract and tubes were spun at 14,500 RPM for 20 minutes at 25°C. Supernatant was collected, pooled, and heat inactivated at 56°C prior to sterile filtration and freezing.

Purified human monocytes were cultured in 10% PDS/RPMI containing 5 units/ml of Genzyme recombinant human MCSF for 5 days before being treated with LDL, oxidized LDL, acetylated LDL (all LDL at 50 µg/ml), lysophosphatidylcholine (Sigma, 37.5 µM), or homocysteine (Sigma, 1mM). After incubation with these reagents for periods ranging from 2 hours up to 3 days, the media was withdrawn and the cells were dissolved in RNA lysis buffer and RNA was prepared as described, above, in Section 6.1.

Lipoproteins For oxidation, human LDL (Sigma) was first diluted to 1 mg/ml with PBS and then dialyzed against PBS at 4°C overnight. LDL was then diluted to 0.3 mg/ml with PBS. CuSO₄·5H₂O was then added to 5µM final concentration, and the solution was incubated in a T flask in a 37°C incubator for 24 hr. LDL solution was then dialyzed at 4°C against 0.15M NaCl/0.3mM EDTA for 2 days with several changes, before being removed and concentrated using an Amicon spin column by spinning for 1 hr. 4000 RPM at 4°C.

For acetylation, 1 ml of 5 mg/ml LDL was added to 1 ml of a saturated solution of NaOAc in a 15 ml tube on ice on a shaker at 4°C. 8 µl of acetic anhydride was added 2 µl at a time over 1 hr. LDL was then dialyzed for 48 hr. against 0.15M NaCl/0.3 mM EDTA at 4°C for 48 hr. with several changes. Final concentrations of derivatized LDL's were determined by comparing to a dilution curve of native LDL analyzed at OD₂₈₀, with 0.15M NaCl/0.3mM EDTA used as diluent in all cases.

6.1.2. ANALYSIS OF PARADIGM MATERIAL

Differential Display:

Removal of DNA: The RNA pellet was resuspended in H₂O and quantified by spectrophotometry at OD₂₆₀. Approximately
5 half of the sample was then treated with DNase I to remove
contaminating chromosomal DNA. RNA was amplified by PCR
using the following procedure. 50 μ l RNA sample (10-20 μ g),
5.7 μ l 10x PCR buffer (Perkin-Elmer/Cetus), 1 μ l RNase
inhibitor (40 units/ μ l) (Boehringer Mannheim, Germany) were
10 mixed together, vortexed, and briefly spun. 2 μ l DNase I (10
units/ μ l) (Boehringer Mannheim) was added to the reaction
which was incubated for 30 min. at 37°C. The total volume
was brought to 200 μ l with DEPC H₂O, extracted once with
phenol/chloroform, once with chloroform, and precipitated by
15 adding 20 μ l 3M NaOAc, pH 4.8, (DEPC-treated), 500 μ l
absolute ETOH and incubating for 1 hour on dry ice or -20°C
overnight. The precipitated sample was centrifuged for 15
min., and the pellet was washed with 70% ETOH. The sample
was re-centrifuged, the remaining liquid was aspirated, and
20 the pellet was resuspended in 100 μ l H₂O. The concentration
of RNA was measured by reading the OD₂₆₀.

First strand cDNA synthesis: For each RNA sample
duplicate reactions were carried out in parallel. 400 ng RNA
25 plus DEPC H₂O in a total volume of 10 μ l were added to 4 μ l
T₁₁XX reverse primer (10 μ M) (Operon). The mixture was
incubated at 70°C for 5 min. to denature the RNA and then
placed at r.t. 26 μ l of reaction mix containing the
following components was added to each denatured RNA/primer
30 sample: 8 μ l 5x First Strand Buffer (Gibco/BRL,
Gaithersburg, MD), 4 μ l 0.1M DTT (Gibco/BRL), 2 μ l RNase
inhibitor (40 units/ μ l) (Boehringer Mannheim), 4 μ l 200 μ M
dNTP mix, 6 μ l H₂O, 2 μ l Superscript reverse transcriptase
(200 units/ μ l) (Gibco/BRL). The reactions were mixed gently
35 and incubated for 30 min. at 42°C. 60 μ l of H₂O (final volume
= 100 μ l) were then added and the samples were denatured for
5 min. at 85°C and stored at -20°C.

PCR reactions: 13 μ l of reaction mix was added to each tube of a 96 well plate on ice. The reaction mix contained 6.4 μ l H₂O, 2 μ l 10x PCR Buffer (Perkin-Elmer), 2 μ l 20 μ M dNTP's, 0.4 μ l ³⁵S dATP (12.5 μ Ci/ μ l; 50 μ Ci total) 5 (Dupont/NEN), 2 μ l forward primer (10 μ M) (Operon), and 0.2 μ l AmpliTaq Polymerase (5 units/ μ l) (Perkin-Elmer). Next, 2 μ l of reverse primer (T₁₁XX, 10 μ M) were added to the side of each tube followed by 5 μ l of cDNA also to the sides of the tubes, which were still on ice. Tubes were capped and mixed, 10 and brought up to 1000 RPM in a centrifuge then returned immediately to ice. The PCR machine (Perkin-Elmer 9600) was programmed for differential display as follows:

	94°C	2 min.
	*94°C	15 sec.
15	*40°C	2 min.
	*ramp 72°C	1 min.
	*72°C	30 sec.
	72°C	5 min.
	4°C	hold

* = X40

20 When the PCR machine reached 94°C, the plate was removed from ice and placed directly into the Perkin-Elmer 9600 PCR machine . Following PCR, 15 μ l of loading dye, containing 80% formamide, 10 mM EDTA, 1 mg/ml xylene cyanol, 25 1 mg/ml bromphenol blue were added. The loading dye and reaction were mixed, incubated at 85°C for 5 min., cooled on ice, centrifuged, and placed on ice. Approximately 4 μ l from each tube were loaded onto a prerun (60V) 6% acrylamide gel. The gel was run at approximately 80V until top dye front was 30 about 1 inch from bottom. The gel was transferred to 3MM paper (Whatman Paper, England) and dried under vacuum. Bands were visualized by autoradiography.

Band isolation and amplification: Differentially expressed 35 bands were excised from the dried gel with a razor blade and placed into a microfuge tube with 100 μ l H₂O and heated at 100°C for 5 min., vortexed, heated again to 100°C for 5 min.,

and vortex again. After cooling, 100 μ l H₂O, 20 μ l 3M NaOAc, 1 μ l glycogen (20 mg/ml), and 500 μ l ethanol were added and chilled. After centrifugation, the pellet was washed and resuspended in 10 μ l H₂O.

- 5 The isolated differentially expressed bands were then amplified by PCR using the following reaction conditions:

	58	μ l	H ₂ O
	10	μ l	10x PCR Buffer
	10	μ l	200 μ M dNTP's
10	10	μ l	10 μ M reverse primer
	10	μ l	10 μ M forward primer
	1.5	μ l	amplified band
	0.5	μ l	AmpliTaq polymerase (5 units/ μ l) (Perkin Elmer)

- 15 PCR was performed using the program described in this Section, above, for differential display. After PCR, glycerol loading dyes were added and samples were loaded onto a 2% preparative TAE/Biogel (Bio101, La Jolla, CA) agarose gel and eluted. Bands were then excised from the gel with a
- 20 razor blade and vortexed for 15 min. at r.t., and purified using the Mermaid kit from Bio101 by adding 3 volumes of Mermaid high salt binding solution and 8 μ l of resuspended glassfog in a microfuge tube. Glassfog was then pelleted, washed 3 times with ethanol wash solution, and then DNA was
- 25 eluted twice in 10 μ l at 50°C.

- Subcloning: The TA cloning kit (Invitrogen, San Diego, CA) was used to subclone the amplified bands. The ligation reaction typically consisted of 4 μ l sterile H₂O, 1 μ l
- 30 ligation buffer, 2 μ l TA cloning vector, 2 μ l PCR product, and 1 μ l T4 DNA ligase. The volume of PCR product can vary, but the total volume of PCR product plus H₂O was always 6 μ l. Ligations (including vector alone) were incubated overnight at 12°C before bacterial transformation. TA cloning kit
- 35 competent bacteria (INV α F': *enda1*, *recA1*, *hsdR17*(r-k, m+k), *supE44*, λ -, *thi-1*, *gyrA*, *relA1*, ϕ 80lacZ α Δ M15 Δ (*lacZYA-argF*), *deoR*+, F') were thawed on ice and 2 μ l of 0.5 M β -

mercaptoethanol were added to each tube. 2 μ l from each ligation were added to each tube of competent cells (50 μ l), mixed without vortexing, and incubated on ice for 30 min. Tubes were then placed in 42°C bath for exactly 30 sec.,
 5 before being returned to ice for 2 min. 450 μ l of SOC media (Sambrook et al., 1989, *supra*) were then added to each tube which were then shaken at 37°C for 1 hr. Bacteria were then pelleted, resuspended in ~200 μ l SOC and plated on Luria broth agar plates containing X-gal and 60 μ g/ml ampicillin
 10 and incubated overnight at 37°C. White colonies were then picked and screened for inserts using PCR.

A master mix containing 2 μ l 10x PCR buffer, 1.6 μ l 2.5 mM dNTP's, 0.1 μ l 25 mM MgCl₂, 0.2 μ l M13 reverse primer (100 ng/ μ l), 0.2 μ l M13 forward primer (100 ng/ μ l), 0.1 μ l
 15 AmpliTaq (Perkin-Elmer), and 15.8 μ l H₂O was made. 40 μ l of the master mix were aliquoted into tubes of a 96 well plate, and whole bacteria were added with a pipette tip prior to PCR. The PCR machine (Perkin-Elmer 9600) was programmed for insert screening as follows:

20	94°C	2 min.
	*94°C	15 sec.
	*47°C	2 min.
	* = X35	*ramp 72°C 30 sec.
		*72°C 30 sec.
25	72°C	10 min.
	4°C	hold

Reaction products were eluted on a 2% agarose gel and compared to vector control. Colonies with vectors containing inserts were purified by streaking onto LB/Amp plates.
 30 Vectors were isolated from such strains and subjected to sequence analysis, using an Applied Biosystems Automated Sequencer (Applied Biosystems, Inc. Seattle, WA).

Northern analysis: Northern analysis was performed to
 35 confirm the differential expression of the genes corresponding to the amplified bands. The probes used to detect mRNA were synthesized as follows: typically 2 μ l

along with a fluorescent ruler before blotting with Hybond-N membrane (Amersham), according to the methods of Sambrook et al., 1989, *supra*, in 20x SSC overnight.

For hybridization, the blot was placed into a roller bottle containing 10 ml of prehybridization solution consisting of 50% formamide and 1x Denhardt's solution, and placed into 65°C incubator for 30 min. The probe was then heated to 95°C, chilled on ice, and added to 10 ml of hybridization solution, consisting of 50% formamide, 1x Denhardt's solution, 10% dextran sulfate, to a final concentration of 10⁶ cpm/ml. The prehybridization solution was then replaced with the probe solution and incubated overnight at 42°C. The following day, the blot was washed three times for 30 min. in 2x SSC/0.1% SDS at room temperature before being covered in plastic wrap and put down for exposure.

RT-PCR Analysis: RT-PCR was performed to detect differentially expressed levels of mRNA from the genes corresponding to amplified bands. First strand synthesis was conducted by mixing 20 µl DNase RNA (~2 µg), 1 µl oligo dT (Operon) (1 µg), and 9.75 µl H₂O. The samples were heated at 70°C for 10 min., and then allowed to cool on ice. 10 µl first strand buffer (Gibco/BRL), 5 µl 0.1M DTT, 1.25 µl 20 mM dNTP's (500 µM final), 1 µl RNasin (40 units/µl) (Boehringer Mannheim), and 2 µl Superscript Reverse Transcriptase (200 units/µl) (Gibco/BRL) were added to the reaction, incubated at 42°C for 1 hr., and then placed at 85°C for 5 min., and stored at -20°C.

PCR was performed on the reverse transcribed samples. Each reaction contained 2 µl 10x PCR buffer, 14.5 µl H₂O, 0.2 µl 20 mM dNTP's (200 µM final), 0.5 µl 20 µM forward primer (0.4 µM final), 0.5 µl 20 µM reverse primer (0.4 µM final), 0.3 µl AmpliTaq polymerase (Perkin-Elmer/Cetus), 2 µl cDNA dilution or positive control (~40 pg). Samples were placed in the PCR 9600 machine at 94°C (hot start), which was programmed as follows:

7. EXAMPLE: IDENTIFICATION OF GENES DIFFERENTIALLY
EXPRESSED IN RESPONSE TO PARADIGM B: IN VIVO MONOCYTES

In an alternative embodiment of the invention, genes differentially expressed in monocytes were detected under
5 highly physiologically relevant, in vivo conditions. According to Paradigm B, human subjects were held in a clinical setting and the fat/cholesterol content of their diets was controlled. Monocytes were isolated at different stages of treatment, and their gene expression pattern was
10 compared to that of control groups.

By use of Paradigm B, the human bcl-2 gene was identified. Its expression decreases in response to the atherogenic conditions of high fat/high cholesterol. The Apo E-/- mouse is the first mouse model of atherosclerosis with
15 pathology similar to that of human plaque development (Plump et al., 1992, Cell 71: 343-353). Serum cholesterol levels in these mice on a chow diet is five times higher than those of control littermates. To address whether the regulation of the mouse bcl-2 gene is also affected by serum cholesterol
20 levels, white blood cells, which include monocytes, from apoE-deficient mice and littermate wild-type controls were purified and mouse bcl-2 mRNA levels were compared using quantitative RT-PCR. By this method, mouse bcl-2 mRNA levels were significantly lower in the apoE-deficient mice relative
25 to the wild-type controls.

The differential expression pattern of the human glutathione peroxidase gene (HUMGPXP1) was also discovered. The differential expression of HUMGPXP1 was initially detected in a preliminary detection system, described, below,
30 in Section 7.1.2. Once HUMGPXP1 sequences were obtained, the gene's differential expression pattern was verified and characterized under the physiologically relevant conditions of Paradigm B. Glutathione peroxidase is known to be involved in the removal of toxic peroxides that form in the
35 course of growth and metabolism under normal aerobic conditions and under oxidative stress. Human plasma glutathione peroxidase gene was originally isolated from a

human placenta cDNA library (Takahashi et al., 1990, J. Biochem. 108: 145-148). It has been shown to be expressed in two human cell lines of the myeloid lineage (Porter et al., 1992, Clinical Science 83: 343-345). Other studies have also
5 linked reduced levels of this enzyme with heart attack risk (Guidi, et al., 1986, J. Clin. Lab Invest. 46: 549-551; Wang et al., 1981, Klin. Wochenschr. 59: 817-818; Kok et al., 1989, J. Am. Med. Assoc. 261: 1161-1164; and Gromadzinska & Sklodowska, 1990, J. Am. Med. Assoc. 263: 949-950).
10 Glutathione peroxidase has not been previously known to be down-regulated in human monocytes under cardiovascular disease conditions, as described herein.

Interestingly, bcl-2 has been recognized as playing a key role in preventing apoptosis, and expression of
15 glutathione peroxidase in the absence of bcl-2 is able to compensate for this loss by preventing apoptosis (Hockenbery et al., 1993, Cell 75: 241-251). These findings regarding bcl-2 and HUMGPXP1, described herein in this section, suggested a novel role for the monocyte in plaque formation
20 which involves apoptosis induction caused by high LDL concentrations inside the cell, or perhaps by oxidative stress in the cell mediated by oxidized LDL.

To confirm this relationship between apoptosis and atherosclerosis, the ability of bcl-2 expression to
25 ameliorate atherosclerosis is tested. Because bcl-2 is normally down-regulated under atherogenic conditions, a transgenic mouse strain is engineered in which the human bcl-2 gene is expressed under the control of the scavenger receptor promoter, which is induced in monocyte foam cells
30 under atherogenic conditions. This transgenic mouse is then crossed with an apoE-deficient atherosclerotic mouse model. The ability of the increased expression of the bcl-2 target gene to ameliorate atherosclerosis is demonstrated by a decrease in initiation and progression of plaque formation
35 observed in the transgenic apoE-deficient mouse.

The identification of the differential expression of these genes, therefore, provides targets for the treatment

and diagnosis of cardiovascular disease. Intervening in the apoptotic pathway through Bcl-2 and glutathione peroxidase, may lead to lesion regression or prevention of plaque formation, or both. Furthermore, the discovery of a connection between the apoptotic pathway and atherosclerosis demonstrates the effectiveness of the methods described herein in identifying the full panoply of gene products that are involved in the atherosclerotic disease process. Furthermore, the down-regulation of bcl-2 and HUMGPXP1 under Paradigm B provides a fingerprint for the study of the effect of excess LDL on monocytes.

7.1. MATERIALS AND METHODS

7.1.1. IN VIVO CHOLESTEROL STUDIES

Patients were held in a clinical setting for a total of 9 weeks during which time their lipid intake was very tightly controlled. There were a total of 3 diets, and each patient was held on each diet for 3 weeks. Patients were healthy young (third decade of life) individuals with no history or symptoms of heart disease or dislipidemias. The 3 diets are described below:

American Heart Association Diet II

fat	25%
cholesterol	80 mg/1000 kCal
polyunsaturated/saturated fat	1.5

Average American Diet

fat	43%
cholesterol	200 mg/1000 kCal
polyunsaturated/saturated fat	0.34

Combination Diet

fat	43%
cholesterol	80 mg/1000 kCal
polyunsaturated/saturated fat	0.34

The 3 diets were isocaloric, and the individual components of each diet may vary with the participant's preference as long as the lipid levels in the diet were maintained.

5 Cell Isolation

At the end of each 3 week diet period, blood was drawn from each patient after a 12 hour period of fasting and monocytes were purified. 50 ml of blood was drawn into 5 evacuated tubes containing 1.4 ml each of citrate phosphate
10 dextrose to prevent coagulation. Blood was pooled into 50 ml tubes and spun at 400g (1250 RPM/Sorvall RC3B) for 15 minutes at 4°C. The upper serum layer (~ 25 ml) was then removed with a pipette and replaced with phosphate buffered saline (PBS) at 4°C. The blood was mixed and then spun at 1850 x g
15 (2680 RPM) for 15 minutes at 4°C. Most of the clear upper layer was removed with a pipette, before the buffy coat at the interface was taken in ~5 ml. The buffy coat was placed into a separate 50 ml tube, and the pipette used to remove it was washed with 20 ml PBS. A small aliquot of these cells
20 was then diluted 1:1000 in PBS and counted under a microscope using a hemacytometer. Red blood cell concentration was then adjusted with PBS to a final concentration of 1.5×10^9 /ml, and 10 ml aliquots were added to Leucoprep Becton Dickinson) tubes for monocyte isolation. Tubes were spun for 25 minutes
25 at 25°C in a Sorvall RT6000 with the brake off. Most of the clear upper layer was discarded, and the turbid layer above the gel was saved and pooled in 50 ml tubes. The volume of each tube was then increased to 50 ml with 25°C PBS, and spun at 1000 RPM (Sorvall RC3B) for 10 minutes at 4°C. The liquid
30 was then discarded, the pellet was resuspended in 50 ml PBS, and spun again. This process was repeated 3 more times. The final cell pellet was then resuspended in 2 ml RNA lysis buffer (Sambrook et al., 1989, supra) and frozen for subsequent RNA isolation as described above in Section 6.1.1.
35 Differential display, Northern analysis, RT-PCR, subcloning, and DNA sequencing were carried out as described, above, in Section 6.1.2.

7.1.2. PRELIMINARY DETECTION SYSTEM

The preliminary detection system described in this section was used to identify sequences that are differentially expressed in a readily assayed, in vitro
5 system. Sequences that showed some homology to those thought to be involved in cardiovascular disease were then used as specific primers or probes, or both, in Paradigm B, wherein the differential expression was ascertained under physiologically relevant conditions, as described in section
10 7.1.1, above.

Cell culture Blood (~100 ml) was drawn from healthy human donors into vacutainer tubes containing heparin (Becton Dickinson). Blood was diluted 1:1 with PD (Phosphate buffered saline (PBS) without Ca or Mg, plus 0.3mM EDTA), and
15 layered onto Ficoll (Lymphocyte Separation Media - Organon Teknicon) as 30 ml of blood/7 ml ficoll in a 50 ml blue-capped Falcon tube, and centrifuged at 2000 RPM for 25 min. at room temperature (r.t.). The buffy coat was removed with a pipette, transferred to another 50 ml tube, diluted to 30
20 ml with PD, and centrifuged at 1200 RPM for 10 min. at r.t. The pellet was resuspended in 30 ml PD and the previous centrifugation step was repeated. The pellet was resuspended in 40 ml RPMI (2mM 1-Glutamine + penicillin/streptomycin), plated onto 4 plates, and incubated at 37°C for 2 hours.
25 Supernatant was removed, and the plates were washed 3x with PBS at 37°C. Plates were finally resuspended in 10 ml each with RPMI/20% human AB serum (Sigma, St. Louis, MO). On day 5, the media was changed and 100 units/ml of human γ -IFN (Genzyme) were added. On day 7, the media was removed and
30 replaced with RPMI/20% human LDL-deficient serum + 100 units/ml of human γ -IFN. Native, oxidized, and acetylated LDL were each added to one plate with the fourth plate serving as control. After the specified incubation time (5 hr. or 24 hr.) the media was removed and the cells were
35 resuspended in 2 ml guanidine isothiocyanate RNA lysis buffer (Sambrook et al., 1989, *supra*). Lysed cells were then syringed with 23 G. needle, layered over 5.7M CsCl, and

centrifuged for 20 hr. at 35K RPM. RNA was isolated according to the method of Sambrook et al., 1989, *supra*.

Lipoproteins were prepared as described, above, in section 6.1.1. Differential display, Northern analysis, RT-PCR, subcloning, and DNA sequencing were carried out as described, above, in Section 6.1.2. For differential display, the primers used were T₁₁CC (reverse) and OPE4 (forward), consisting of 5'GTGACATGCC3'. For RT-PCR, the first strand cDNA was primed with T₁₁CC, and PCR reactions were carried out with rfhma15 primers (for- catgcctgtagaaaaaggtt/rev-cttcatagaatctaagccta), and mouse γ actin primers (for-cctgatagatgggcactgtgt/rev-gaacacggcattgtcactaact). (SEQ ID NO:1) (SEQ ID NO:2) (SEQ ID NO:3) (SEQ ID NO:4)

7.1.3. TRANSGENIC ApoE-DEFICIENT MOUSE EXPRESSING HUMAN bcl-2

Transgenic mice bearing a construct with the mouse scavenger receptor regulatory element (5kb) (M. Freeman, et al., 1995, unpublished results) driving expression of the human bcl-2 gene (hbcl-2) were produced. The scavenger receptor regulatory element (ScR) is known to activate reporter gene expression in peritoneal macrophages in transgenic mice (M. Freeman, 1995, unpublished results). This 5 kb fragment is linked to the human bcl-2 cDNA (Cleary, et al., 1986, *supra*) via a NotI restriction site. Human growth hormone (hGH) sequences (Mayo, et al., 1983, Nature 306: 86-88) are then ligated onto the 3' end of this construct through filled-in BamHI and EcoRV sites to provide message stability. This construct is then digested with XhoI and the 9 kb ScR-hbcl2-hGH sequences are purified away from vector sequences. Another plasmid sample is digested with KpnI to yield a fragment with only 1.5kb of scavenger receptor regulatory sequences which provide a lower level of expression. These fragments are then injected independently into mouse embryos derived from the FVB and C57BL/6 mouse strains according to standard protocols (Hogan, et al., Manipulating the Mouse Embryo, 1994, Cold Spring Harbor

Laboratory Press). Following birth, tail sections are cut from mice derived from injected embryos and analyzed for the presence of transgene sequences using hbcl-2 sequences as probes on Southern blots.

5 Transgenic mice bearing the ScR-hbcl2-hGH construct are then bred to wild-type mice of the same respective strain, and then the offspring are backcrossed to produce homozygous lines of mice. These mice are then bred to apoE-deficient mice. Offspring are analyzed for presence of the
10 ScR-hbcl2-hGH by preparing tail sections and probing with hbcl-2 sequences on Southern blots. Offspring are then analyzed for lesion formation and progression according to the methods of Plump, et al., 1992, *supra*.

Differential display analysis was carried out on monocyte RNA derived from the blood of patients whose serum cholesterol levels were manipulated through fat/cholesterol intake in their diets. A band designated band #14 which was present in the low dietary fat/low serum cholesterol conditions and goes away in the high dietary fat/high serum cholesterol conditions. When a radioactively labeled probe was prepared from band #14 and hybridized with a Northern blot prepared from RNA from the same patient, an 8 kb band was seen which was present in low serum cholesterol and disappeared in high serum cholesterol conditions. When band #14 sequences were subcloned, sequenced, and compared with the sequence database a 98% (203/207 bp) sequence similarity with the human bcl-2 gene (Cleary et al., 1986, Cell 47, 19-28) was obtained, indicating that band #14 is bcl-2.

analysis was carried out, and regulated bands were isolated and characterized. The DNA sequence was determined from a number of independent subclones of amplified sequences of one such regulated band designated band 15. Using the BLAST
5 program (Altschul, et al., 1990, J. Mol. Biol. 215: 403-410), a 176/177 (99%) sequence similarity was found between band 15 a sequence for human plasma glutathione peroxidase exon 1 (HUMGPXP1). This sequence occurs upstream of the reported transcription start site. Nonetheless, RT-PCR analysis
10 confirmed that the band 15 sequences are in fact within the same transcription unit as sequences downstream of the reported transcription start site.

Based on this preliminary result, the gene expression pattern of glutathione peroxidase (HUMGPXP1) was further
15 analyzed for verification and characterization in physiologically relevant samples according to Paradigm B. Monocytes derived from human blood under atherogenic conditions (high serum cholesterol) and healthy conditions (low serum cholesterol) were examined with RT-PCR. There
20 appeared to be 2-3 fold less cDNA amplified by the HUMGPXP1 primers from the high fat/cholesterol monocytes than in the low fat/cholesterol monocytes, while the actin control bands are the same.

White blood cells, which include monocytes, from apoE-
25 deficient mice and littermate wild-type controls were purified, and mouse bcl-2 mRNA levels were compared using quantitative RT-PCR. By this method, mouse bcl-2 mRNA levels were significantly lower in the apoE-deficient mice relative to the wild-type controls.

30 These results demonstrate that bcl-2 is an excellent target gene for intervening in lesion formation and development. It was previously known that, under normal conditions, bcl-2 expression prevents apoptosis. The observed down-regulation of bcl-2 caused by atherogenic
35 conditions, therefore, provides an explanation of how such atherogenic conditions may lead to plaque formation. By down-regulating the normally protective bcl-2 gene, high

serum cholesterol triggers a series of events, entailing the induction of the apoptotic pathway, which results in programmed cell death, which in turn causes an inflammatory response and subsequent plaque formation.

5 This model may be tested by counteracting the observed down-regulation of bcl-2. The human bcl-2 gene is placed in the ScR-hbcl2-hGH construct in which it is transcribed by a promoter that is activated in monocyte foam cells under atherogenic conditions. This construct is then introduced
10 into an apoE-deficient mouse that otherwise serves as a model for atherosclerosis. The effect of bcl-2 expression on atherosclerosis is evidenced by the reduction in plaque initiation and development in the apoE-deficient mice bearing the construct. Amelioration of atherosclerosis may,
15 therefore, be accomplished by such intervention in the down-regulation of the bcl-2 target gene.

8. EXAMPLE: IDENTIFICATION OF GENES DIFFERENTIALLY EXPRESSED IN RESPONSE TO PARADIGM C: IL-1 INDUCTION OF ENDOTHELIAL CELLS

20 According to the invention, differential display was used to detect four novel genes that are differentially expressed in endothelial cells that were treated in vitro with IL-1. Three of these genes, rchd024, rchd032, and
25 rchd036, are not homologous to any known gene. The fourth gene, rchd005, is 70% homologous to a cloned shark gene called bumetanide-sensitive Na-K-Cl cotransport protein. A human homolog of this gene has been reported, but the sequence has not yet been published (Xu et al., 1994, Proc.
30 Natl. Acad. Sci. USA 91: 2201-2205).

The discovery of the up-regulation of these four genes provides a fingerprint profile of IL-1 induced endothelial cells. This fingerprint profile can be used in the treatment and diagnosis of cardiovascular diseases, including but not limited to atherosclerosis, ischemia/reperfusion, hypertension, restenosis, and arterial inflammation.

8.1. MATERIALS AND METHODS

Primary cultures of HUVEC's were established from normal term umbilical cords as described (In Progress in Hemostasis and Thrombosis, Vol. 3, P. Spaet, editor, Grune & Stratton Inc., New York, 1-28). Cells were grown in 20% fetal calf serum complete media (Luscinskas, et al., 1989, J. Immunol. 142: 2257-2263) and passaged 1-3 times before activation.

For activation, cells were cultured with 10 units/ml of human IL-1 β for 1 or 6 hr. before lysis in guanidinium isothiocyanate RNA lysis buffer (Sambrook et al., 1989, supra). Lysed cells were then syringed with a 23 G. needle, layered over 5.7M CsCl, and centrifuged for 20 hr. at 35K.

Alternatively, cells were induced in the presence of 100 μ M lysophosphatidylcholine, or 50 μ g/ml oxidized human LDL (Sigma) for periods of 1 or 6 hr. RNA was isolated as described, above, in Section 6.1. Differential display, Northern analysis, RT-PCR, subcloning, and DNA sequencing were carried out as described, above, in Section 6.1.2, except that Northern blot hybridizations were carried out as follows: for pre-hybridization, the blot was placed into roller bottle containing 10 ml of rapid-hyb solution (Amersham), and placed into 65°C incubator for at least 1 hr. For hybridization, 1x10⁷ cpm of the probe was then heated to 95°C, chilled on ice, and added to 10 ml of rapid-hyb solution. The prehybridization solution was then replaced with probe solution and incubated for 3 hr at 65°C. The following day, the blot was washed once for 20 min. at r.t. in 2x SSC/0.1% SDS and twice for 15 min. at 65°C in 0.1x SSC/0.1% SDS before being covered in plastic wrap and put down for exposure.

Chromosomal locations were determined according to the method described in Section 6.1.3, above. For rchd024, the primers used were for-cccatagactaggctcatag^(SEQ ID NO:5), and rev-tttaaagagaaattcaa^(SEQ ID NO:6)tc.

8.2. RESULTS

HUVEC's were activated with 10 units/ml IL-1 β for 1 or 6 hours and compared to resting HUVEC's using differential display. A band designated rchd005 was present in lanes 11 and 12 (IL-1, 6 hr.) but not in lanes 9 and 10 (control), or lanes 7 and 8 (IL-1, 1 hr.). This band, rchd005, was isolated and subcloned and sequenced. When a probe prepared from this band was used to screen a Northern blot, expression was seen at 6 hr., but not at 1 hr. or in the control. However, when this same probe was hybridized to a Northern blot prepared from shear stressed RNA, according to Paradigm D described in Section 9, below, a different pattern of up-regulation was also seen. Expression was up at 1 hr. and then nearly disappeared by 6 hr. Amplified rchd005 DNA was subcloned and sequenced. Sequence analysis revealed an approximately 360 bp insert (FIG.1) with 70% sequence similarity to a cloned shark gene called bumetanide-sensitive Na-K-Cl cotransport protein.

Another IL-1 inducible differential display band was designated rchd024. Northern analysis on IL-1 up-regulated RNA revealed a 10 kb rchd024 message present at 6 hr. that also showed a low level of up-regulation under shear stress at 6 hr. The DNA sequence of rchd024 was obtained from subclones of amplified DNA (FIG.2). Database searching revealed no significant sequence similarities. A PCR amplification experiment determined that the rchd024 gene is located on human chromosome 4.

Band rchd032 was isolated on the basis of its differentially increased expression after 6 hr. treatment with IL-1, which was confirmed by RT-PCR analysis. Amplified rchd032 sequences were subcloned and sequenced (FIG.3). No significant homology to any known gene was found.

Band rchd036 was also isolated on the basis of its differential expression 6 hr. after IL-1 treatment. Northern analysis revealed an 8 kb band which was up-regulated 6 hr. after IL-1 treatment. Another Northern analysis was performed testing rchd036 under the shear stress condition of

Paradigm D, which are described in the example in Section 9, below. Interestingly, rchd036 is not induced by shear stress, as indicated by the lack of any band after either 1 hr. or 6 hr. of treatment. This result provides an example
5 of an IL-1-inducible endothelial cell gene that is not regulated by shear stress, indicating that these induction pathways can be separated, and may provide for drugs with greater specificity for the treatment of inflammation and atherosclerosis. The DNA sequence was obtained from
10 subclones of amplified DNA (FIG.4), and a search of the database revealed no sequence similarities. A PCR amplification experiment determined that the rchd036 gene is located on human chromosome 15.

15 9. EXAMPLE: IDENTIFICATION OF GENES DIFFERENTIALLY EXPRESSED IN RESPONSE TO PARADIGM D: ENDOTHELIAL CELL SHEAR STRESS

According to the invention, differential display was used to detect genes that are differentially expressed in
20 endothelial cells that were subjected to fluid shear stress in vitro. Shear stress is thought to be responsible for the prevalence of atherosclerotic lesions in areas of unusual circulatory flow. Using the method of Paradigm D, four bands with novel DNA sequences were identified.

25 rchd502 is homologous to rat matrin F/G mRNA sequence (Hakes, et al., 1991, Proc. Natl. Acad. Sci. U.S.A. 88:6186-6190). This rat gene has been shown to encode a protein which functions as a prostaglandin transporter, and has been designated PGT (Kanai et al., 1995, Science 268: 866-869).
30 In fact, the sequences in rchd502 encode the homologous twelve transmembrane domains found in the PGT gene. Furthermore, rchd502 was demonstrated to be up-regulated by shear-stress but not by IL-1. It therefore provides an excellent novel tool for diagnosis and treatment of
35 cardiovascular disease.

The complete sequence of the rchd523 gene reveals that it encodes a novel G protein-coupled receptor protein,

consisting of 375 amino acids and seven transmembrane domains. At the amino acid level, rchd523 is 40% indentical to the Angiotensin II receptor. The discovery of such a novel protein is particularly useful in designing treatments
5 as well as diagnostic and monitoring systems for cardiovascular disease. In carrying out signal transduction, G proteins play an important early role in the pathways that cause changes in cellular physiology. The rchd523 gene product, therefore, provides an excellent target for
10 intervention in the treatment of cardiovascular disease.

Furthermore, as transmembrane proteins, the rchd502, rchd523, and rchd528 gene products can be readily accessed or detected on the endothelial cell surface by other compounds. They provide, therefore, excellent targets for detection of cardiovascular disease states in diagnostic systems, as well as in the monitoring of the efficacy of compounds in clinical trials. Furthermore, the extracellular domains of these four gene products provide especially efficient screening systems for identifying compounds that bind to them. Such compounds, can be useful in treating cardiovascular disease by modulating the activity of the transmembrane gene products.

protein-4/Transforming growth factor- β (TGF- β). As described in detail in Section 16, below, the 235 amino acid rchd534 protein is encoded by the shorter of two spliceoforms of the rchd534 gene.

- 5 Also using the method of Paradigm D, the previously identified human prostaglandin endoperoxide synthase type II, also known as cyclooxygenase II (COX II), was identified (band rchd505). This gene was previously known to be involved in inflammation, and to be up-regulated by IL-1
- 10 (Jones et al., 1993, J. Biol. Chem. 268: 9049-9054), but its up-regulation by shear stress was previously unknown. This result confirmed the general effectiveness of the techniques used according to the invention in the detection of genes involved cardiovascular disease.
- 15 The sequence of another up-regulated gene, designated as rchd530, was shown to be identical to the previously identified human manganese superoxide dismutase gene (MnSOD). The up-regulation of MnSOD under shear stress was not previously known.
- 20 The up-regulation of these six genes in shear stressed endothelial cells provides a fingerprint for the study of cardiovascular diseases, including but not limited to atherosclerosis, ischemia/reperfusion, hypertension, and restenosis. The fact that one of these genes, rchd502, is
- 25 not up-regulated under Paradigm C (IL-1 induction) provides an extremely useful means of distinguishing and targeting physiological phenomena specific to shear stress.

- The importance of the induction of these genes in endothelial cells under disease conditions was further
- 30 analyzed by testing the effect of estrogen on their expression. Studies in postmenopausal women on estrogen replacement therapy and in animal models have demonstrated that estrogen has an atheroprotective effect in reducing incidence of coronary artery disease (Gura, T., 1995, Science
- 35 269:771-773). While these studies demonstrate that estrogen has an effect in the liver in reducing LDL levels and increasing HDL levels, these lipoprotein changes are not

thought to be responsible for all of the cardioprotective effects of estrogen.

The identification of target genes that are differentially expressed under certain disease conditions provides for further analysis of the effect of estrogen on cardiovascular disease. The effect of estrogen on target gene expression in endothelial cells was, therefore, compared to particular paradigm expression patterns. Specifically, given that the estrogen receptor is a transcription factor (Kumar and Chambon, Cell 55:145-156, 1988), genes that are induced by shear stress were examined for regulation by estrogen in HUVEC's. In addition to estrogen, treatment with estrogen receptor agonists/antagonists tamoxifen (Grainger et al., Nature Medicine 1:1067-1073, 1995) and raloxifene (Black et al., J. Clin. Invest. 93:63-69, 1994), which also have been reported to have cardioprotective effects, were examined. The results demonstrate that rchd528, which is up-regulated by shear stress, is also up-regulated by estrogen, and suggest that shear stress and estrogen may play similar roles in cardiovascular disease.

9.1. MATERIALS AND METHODS

Primary cultures of HUVEC's were established from normal term umbilical cords as described (In Progress in Hemostasis and Thrombosis, Vol. 3, P. Spaet, editor, Grune & Stratton Inc., New York, 1-28). Cells were grown in 20% fetal calf serum complete media (Luscinskas et al., 1989, J. Immunol. 142: 2257-2263) and passaged 1-3 times before shear stress induction.

For induction, second passage HUVEC's were plated on tissue culture-treated polystyrene and subjected to 10 dyn/cm² laminar flow for 1 and 6 hr. as described (Nagel et al., 1994, J. Clin. Invest. 94: 885-891) or 3-10 dyn/cm² turbulent flow as previously described (Davies et al., 1986 Proc. Natl. Acad. Sci. U.S.A. 83: 2114-2117).

To examine the effect of estrogen on target gene expression, HUVEC's were cultured in serum free endothelial

1989, *supra*). Plaques from the libraries that were detected by the probes were isolated and the cDNA insert within the phage vector was sequenced.

Determination of chromosomal location was carried out according to the method described in Section 6.1.3, above.

Q The primers used for rchd523 were (for-atgccgtgtgggtagtc) (SEQ ID NO: 7)
Q and (rev-attttatgggaagggtttttaca) (SEQ ID NO: 8); and for rchd534 were (for-
a cttttctgcgtctcccat) (SEQ ID NO: 9) and (rev-agacatcagaaactccaacc) (SEQ ID NO: 10).

Northern blot analysis of RNA extracted from various human organs and tissues was performed using commercially available pre-blotted filters (Clontech, Palo Alto, CA).

9.2. RESULTS

HUVEC's were subjected to laminar shear stress for 15 or 6 hr. and compared to static control cells in differential display. A band (rchd502) was identified which was found in lanes 5,6 (6 hr.) but not in lanes 1,2 (control). This band was excised, amplified, and sequenced. Northern analysis using amplified rchd502 sequences revealed a 4.5 kb band that 20 is up-regulated at 6 hr. compared to controls. When rchd502 probe was hybridized to a Northern blot prepared from IL-1 induced endothelial cells, up-regulation of a 4.5 kb band was not seen. This result provides the first example of a shear stress-inducible endothelial cell gene that is not regulated 25 by IL-1, indicating that these induction pathways can be separated, and may provide for drugs with greater specificity for the treatment of inflammation and atherosclerosis. The sequence of the amplified region of rchd502 was used to design probes for cloning the entire gene.

30 Both 5' and 3' RACE reactions were carried out to obtain a 2.2kb cDNA containing the entire coding sequence of the rchd502 gene. Based on the sequence information from RACE, a phage clone was isolated from a human pancreas library which contains all but the first 200 base pairs of 35 the rchd502 coding region. This clone was designated pFCHD502SF. The remaining 200 base pairs were obtained through amplification from a human lung library by PCR with

specific primers. A fragment comprising base pairs 1-265 of the rchd502 gene was subcloned into the TA cloning vector to produce plasmid pFCHD502SJ. Thus, rchd502 is represented by two subclones, pFCHD502SJ comprising base pairs 1-265, and
5 pFCHD502SF comprising base pairs 201 through the 3' end of the coding region, including 3' untranslated sequence.

The complete sequence encompassing the entire coding region is shown in FIG.5. rchd502 shows strong homology (81.4%) to the rat PGT gene, which encodes a prostaglandin
10 transporter (Kanai et al., 1995, *supra*). It contains twelve transmembrane (TM) domains. The approximate bounds of each of the twelve TM domains are as follows:

TM1: about amino acid 31 to about amino acid 52.

TM2: about amino acid 68 to about amino acid 89.

15 TM3: about amino acid 102 to about amino acid 121.

TM4: about amino acid 173 to about amino acid 194.

TM5: about amino acid 206 to about amino acid 227.

TM6: about amino acid 259 to about amino acid 280.

TM7: about amino acid 315 to about amino acid 337.

20 TM8: about amino acid 366 to about amino acid 385.

TM9: about amino acid 403 to about amino acid 423.

TM10: about amino acid 510 to about amino acid 530.

TM11: about amino acid 555 to about amino acid 575.

TM12: about amino acid 607 to about amino acid 627.

25 Shear stress band rchd505 decreased 1 hr. and 6 hr. after shear stress, as compared to untreated control cells. Northern analysis revealed differential expression except that rchd505 was up-regulated after 1 hr. and 6 hr. shear stress treatment. This same band was similarly up-regulated
30 in cells treated with IL-1 according to Paradigm C. Sequence analysis revealed that rchd505 is the previously characterized human endoperoxide synthase type II, also known as cyclooxygenase II (COX II).

rchd523 was detected under differential display as a
35 band up-regulated after 1 hr. and 6 hr. shear stress treatment. The 6 hr. up-regulation of rchd523 was confirmed by RT-PCR. Amplified rchd523 sequences were subcloned, and

an isolate was sequenced and designated pRCHD523. The RACE procedure was used to obtain a 2.5 kb cDNA containing the entire coding sequence of the rchd523 gene. The cDNA isolate containing the complete coding sequence of rchd523 is
5 designated pFCHD523. The DNA sequence comprising the complete coding region of the rchd523 gene is shown in FIG.6. Sequence analysis revealed that the rchd523 gene product encodes a novel G protein-coupled receptor, consisting of 375 amino acids and seven transmembrane domains. At the amino
10 acid level, rchd523 is 40% indentical to the Angiotensin II receptor. A PCR amplification experiment determined that the rchd523 gene is located on human chromosome 7.

rchd528 was also detected as an up-regulated band after 1 hr. and 6 hr. shear stress treatment. This result
15 was confirmed by Northern analysis in which probes of rchd528 amplified sequence detected an approximately 8 kb message that was up-regulated moderately after 1 hr., and up-regulated very strongly after 6 hr. The amplified sequences were subcloned and sequenced. This sequence information was
20 used for initial probing of a cDNA library to isolate the rchd528 gene.

The amplified sequence was used for initial probing of a shear stressed HUVEC cDNA library to isolate a partial clone of rchd528. The RACE procedure was then used in
25 combination with probing a human heart cDNA library and PCR amplification to obtain overlapping clones encompassing the entire rchd528 coding region. The complete coding region of the rchd528 gene is contained in the following three plasmids each containing a segment of the rchd528 gene cloned into
30 pBluescript: pFCHD528A, comprising nucleotides 1-1200; pFCHD528B, comprising nucleotides 237-2982; and pFCHD528C, comprising nucleotides 2982 through the 3' end of the coding region. The DNA sequence comprising the complete coding region of the rchd528 gene is shown in FIG.7.

35 Based on homology to a number of different proteins, the rchd528 gene product was shown to contain an extracellular domain comprising the epidermal growth factor

(EGF) repeat motif. The approximate bounds of the EGF repeat are from about amino acid 1089 to about amino acid 1122.

5 transmembrane domain extending from about amino acid 1348 to about amino acid 1370. In addition, there is an asparagine hydroxylation site consensus sequence from about amino acid 1140 to about amino acid 1151. Northern blot analysis of mRNA isolated from a variety of human organs and tissues
10 revealed that rchd528 is very highly expressed in the heart.

15 A band designated rchd530 corresponded to a sequence strongly up-regulated in HUVECs after six hours of shear stress. This up-regulation is greater for laminar shear stress than for turbulent shear stress. Sequence analysis revealed that rchd530 is identical to human
20 manganese superoxide dismutase (MnSOD). The induction of MnSOD by shear stress was not previously known. MnSOD was also demonstrated to be induced by six hours of treatment with IL-1.

35 An initial comparison with sequences in the database revealed no homologies between rchd534 and any known DNA sequences. A subsequently performed search revealed that

oxidative potential may be measured by performing differential display on the monocytes of patients undergoing clinical tests.

5 10.1. TREATMENT OF PATIENTS AND CELL ISOLATION

Test patients may be administered compounds suspected of having anti-oxidant activity. Control patients may be given a placebo.

Blood may be drawn from each patient after a 12 hour
10 period of fasting and monocytes may be purified as described, above, in Section 7.1.1. RNA may be isolated as described in Section 6.1.1, above.

10.2. ANALYSIS OF SAMPLES

15 RNA may be subjected to differential display analysis as described in Section 6.1.2, above. A decrease in the physiological response state of the monocytes is indicated by a decreased intensity of those bands that were up-regulated by oxidized LDL under Paradigm A, and an increased intensity
20 of those bands that were down-regulated by oxidized LDL under Paradigm A, as described in Section 6.2, above.

11. EXAMPLE: IMAGING OF A CARDIOVASCULAR DISEASE CONDITION

According to the invention, differentially expressed
25 gene products which are localized on the surface of affected tissue may be used as markers for imaging the diseased or damaged tissue. Conjugated antibodies that are specific to the differentially expressed gene product may be administered to a patient or a test animal intravenously. This method
30 provides the advantage of allowing the diseased or damaged tissue to be visualized non-invasively.

For the purposes of illustration, this method is described in detail for the rchd523 gene product. The principles and techniques can be applied to any identified
35 transmembrane target gene product, including, for example, the rchd502 and rchd528 gene products.

11.1. MONOCLONAL CONJUGATED ANTIBODIES

The differentially expressed surface gene product, such as the rchd523 gene product, is expressed in a recombinant host and purified using methods described in Section 5.4.2, above. Preferably, a protein fragment comprising one or more of the extracellular domains of the rchd523 product is produced. Once purified, it is be used to produce F(ab')₂ or Fab fragments, as described in Section 5.4.3, above. These fragments are then labelled with technetium-99m (^{99m}Tc) using a conjugated metal chelator, such as DTPA as described in section 5.8.3, above.

11.2. ADMINISTRATION AND DETECTION OF IMAGING AGENTS

Labeled MAb may be administered intravenously to a patient being diagnosed for atherosclerosis, restenosis, or ischemia/reperfusion. Sufficient time is allowed for the detectably-labeled antibody to localize at the diseased or damaged tissue site (or sites), and bind to the rchd523 gene product. The signal generated by the label is detected by a photoscanning device. The detected signal is then converted to an image of the tissue, revealing cells, such as endothelial cells, in which rchd523 gene expression is up-regulated.

12. EXAMPLE: SCREENING FOR LIGANDS OF THE rchd523 GENE PRODUCT AND ANTAGONISTS OF rchd523 GENE PRODUCT-LIGAND INTERACTION

The rchd523 gene product is a member of the G protein-coupled receptor protein family, containing multiple transmembrane domains. The receptor binding activity of this protein family is detected by assaying for Ca²⁺ mobility through the membrane of cells in which the receptor gene is expressed. This assay, described below, is used to identify ligands that bind to the rchd523 gene product receptor. Establishing this ligand-receptor activity then provides for a screen in which antagonists of the ligand-receptor interaction are identified. An antagonist is detected by its

ability to inhibit the Ca^{2+} mobility induced by ligand-receptor binding. Such antagonists, therefore, provide compounds that are useful in the treatment of cardiovascular disease, by counteracting the activity of the product of this
5 target gene which is up-regulated in the disease state.

Binding of ligand to the rchd523 gene product is measured as follows. The cDNA containing the entire coding region of the rchd523 gene is removed from pFCHD523 and placed under the control of a promoter that is highly
10 expressed in mammalian cells in an appropriate expression vector. The resulting construct is transfected into myeloma cells, which are then loaded with FURA-2 or INDO-1 by standard techniques. Ligands are added to the cell culture to test their ability to bind to the rchd523 receptor in a
15 manner that triggers signal transduction, as measured by Ca^{2+} mobilization across the cell membrane. Mobilization of Ca^{2+} induced by ligand is measured by fluorescence spectroscopy as described in Grynkiewicz et al., 1985, *J. Biol. Chem.* 260:3440. Ligands that react with the target gene product
20 receptor domain are identified by their ability to produce a fluorescent signal. Their receptor binding activities are quantified and compared by measuring the level of fluorescence produced over background.

Candidate antagonists are then screened for their
25 ability to interfere with ligand-receptor binding. Myeloma transfectants expressing rchd523 gene product are treated with ligand alone, and ligand in the presence of candidate antagonist. Candidate antagonists that cause a reduction in the fluorescence signal are designated antagonists of the
30 ligand-rchd523 receptor interaction.

13. POLYCLONAL ANTIBODIES TO TARGET GENE PEPTIDE SEQUENCES

Peptide sequences corresponding to the indicated amino sequences of cDNAs were selected and submitted to Research
35 Genetics (Huntsville, AL) for synthesis and antibody production. Peptides were modified as described (Tam, J.P., 1988, *Proc. Natl. Acad. Sci. USA* 85: 5409-5413; Tam, J.P.,

and Zavala, F., 1989, J. Immunol. Methods 124: 53-61; Tam, J.P., and Lu, Y.A., 1989, Proc. Natl. Acad. Sci. USA 86: 9084-9088), emulsified in an equal volume of Freund's adjuvant and injected into rabbits at 3 to 4 subcutaneous 5 dorsal sites for a total volume of 1.0 ml (0.5 mg peptide) per immunization. The animals were boosted after 2 and 6 weeks and bled at weeks 4, 8, and 10. The blood was allowed to clot and serum was collected by centrifugation.

10

15

20

25

30

35

030428-030430

The peptides used are summarized below:

rchd502 Protein

	<u>Peptide</u>	<u>Amino Acids #'s</u>	<u>Sequence</u>
5			
α	fchd502.1	294-308	DEARKLEEAKSRGSL (SEQ ID NO: 11)
α	fchd502.2	435-449	SSIHPQSPACRRDCS (SEQ ID NO: 12)
α	fchd502.3	627-640	RVKKNKEYNVQKAA (SEQ ID NO: 13)

10

rchd523 Protein

α	fchd523.1	243-258	RAHRHRGLRPRRQKAL (SEQ ID NO: 14)
α	fchd523.2	360-372	IPDSTEQSDVRFSL (SEQ ID NO: 15)

15

rchd528 Protein

α	fchd528.1	1393-1410	SPYAEYPKNPRSQEWGRE (SEQ ID NO: 16)
α	fchd528.2	1467-1481	NPSFISDESRRRDYF (SEQ ID NO: 17)

20

rchd534 and rchd534-long Proteins

		<u>rchd534</u>	<u>rchd534-long</u>	
α	fchd534.1	54-69	-	EFSDASMSPDATKPSH (SEQ ID NO: 18)
α25	fchd534.2	112-125	373-386	LEQRSESVRRTRSK (SEQ ID NO: 19)
α	fchd534.3	182-197	443-458	RSGLQHAPEPDAADGP (SEQ ID NO: 20)

14. **LOCALIZATION OF NOVEL GENES BY *IN SITU* HYBRIDIZATION**

30

The expression of two target genes, rchd502 and rchd528, was examined by *in situ* hybridization. The expression was detected in human carotidendarterectomy samples, i.e., human cardiovascular tissue in a diseased state, taken from a living patient suffering from

35

cardiovascular disease. The expression pattern for each gene was observed to be similar to the pattern detected for the positive control, which is known to be constitutively

Kodak Dektol developer. Slides were counterstained with Haemotoxylin and Eosin and photographed. Controls for the *in situ* hybridization experiments included the use of a sense probe which showed no signal above background levels.

5

14.2 Results

The rchd502 and rchd528 genes each displayed a similar expression patter to the positive control signal from von Willebrand factor, a constitutively expressed endothelial
10 cell marker. Signal was detected for both rchd502 and rchd528 over most endothelial cells lining the luminal surface of the carotid artery, as was also observed for von Willebrand factor. None of the three genes examined showed expression in any other cell type present in the tissue,
15 including smooth muscle cells and macrophages.

15. EXAMPLE: THE RCHD534 AND FCHD540 GENE PRODUCTS INTERACT

The amino acid sequence of the novel rchd534 protein, and the cDNA that encodes it, are described in detail Section
20 9, above. The novel rchd534 gene shares homology with the *Drosophila* Mad gene. The novel fchd540 gene (described in co-pending Application No. 08/799,910, filed February 13, 1997, which is incorporated by reference in its entirety herein) is another gene that is up-regulated in endothelial
25 cells by shear stress. The DNA and encoded amino acid sequence of the fchd540 gene is shown in FIG.10. The fchd540 gene was deposited in the American Type Culture Collection (ATCC) in microorganism pFCHD540 on February 7, 1996 and assigned the ATCC Accession No. 69984. The fchd540 gene also
30 shares homology with the *Drosophila* Mad gene. The fchd540 protein has both an MH1 domain and an MH2 domain. Mad genes have been shown to play a role in the TGF- β signalling pathway (Sekelsky et al., 1995, Genetics 139: 1347-1358; Chen et al., 1996, Nature 383: 691-696; Serra, et al., 1996,
35 Nature Medicine 2: 390-391). TGF- β signalling is considered to be beneficial to atherosclerosis and restenosis (Border et al., 1995, Nature Medicine 1: 1000; Grainger, et al., 1995,

Nature Medicine 1: 1067-1073; Kojima, et al., 1991, J. Cell Biol. 113: 1439-1445; Nikol, et al., 1992, J. Clin. Invest. 90: 1582-1592).

The data described below demonstrate that the rchd534 and fchd540 proteins interact with one another; and this interaction may lead to the inhibition of TGF- β signalling. Furthermore, the expression of these two genes, as described below, is specific to endothelial cells. Because these two genes 1) are both expressed specifically in endothelial cells, 2) are both up-regulated in endothelial cells under certain conditions, 3) encode MAD proteins that interact with one another in endothelial cells, and 4) inhibit TGF- β signalling (which is considered to be beneficial to atherosclerosis), rchd534 and fchd540 proteins are attractive targets for therapeutic intervention in cardiovascular disease. In particular, treatment regimens that inhibit the interaction or activity of the rchd534 and fchd540 proteins can be beneficial for the treatment cardiovascular disease.

Further analyses demonstrated that the rchd534 protein interacts with itself to form a homodimer. Thus, treatment regimens that inhibit the interaction of the rchd534 protein with itself can be beneficial for the treatment cardiovascular disease.

In addition, the analyses described below demonstrated novel interactions of both the rchd534 and fchd540 proteins with other proteins known to be involved in the TGF- β signalling pathway. The protein members of the TGF- β signalling pathway tested included MADR1 (Hoodless et al., 1996, Cell 85:489-500), MADR2 (Eppert et al., 1996, Cell 86: 543-552), DPC4 (Raftery et al., 1988, Genetics 139: 241-254), T β RI, TSR1, ActRIb, ALK3, and ALK6 (Wieser et al., 1995, EMBO J. 14: 2199-2208). For example, the rchd534 protein interacts strongly in endothelial cells with MADR1, MADR2, DPC4, and weakly in 293 (human embryonic kidney) cells with activated forms of receptors T β RI and ActRI. The fchd540 protein interacts strongly in 293 cells with activated forms of receptors T β RI and ALK6.

864020 9246060

In the absence of transfected rchd543 and fchd540 genes, transfected MADR1 or transfected MADR2 mediated a 20-fold induction of a TGF- β inducible promoter in BAECs. Co-expression of either transfected rchd534 or transfected fchd540 in this system eliminated the induction, and also prevented the localization of MADR2 in the nucleus in response to TGF- β signalling. Therefore, treatment regimens that inhibit the interaction of the rchd534 and fchd540 proteins with other proteins involved in the TGF- β pathway also can be beneficial for the treatment cardiovascular of disease. As described above, the expression of rchd534 and fchd540 is specific, within arterial tissue, to endothelial cells. Accordingly, the rchd534 and fchd540 genes may be targets for intervention in a variety of inflammatory and fibroproliferative disorders that involve endothelial cells, including, but not limited to, cancer, angiogenesis, inflammation, and fibrosis.

15.1 MATERIALS AND METHODS

20 15.1.1. YEAST STRAINS, MEDIA, AND MICROBIOLOGICAL TECHNIQUES

Standard yeast media including synthetic complete medium lacking L-leucine, L-tryptophan, and L-histidine were prepared and yeast genetic manipulations were performed as described (Sherman, 1991, Meth. Enzymol., 194:3-21). Yeast transformations were performed using standard protocols (Gietz et al., 1992, Nucleic Acids Res., 20:1425. Ito et al., 1983, J. Bacteriol., 153:163-168). Plasmid DNAs were isolated from yeast strains by a standard method (Hoffman and Winston, 1987, Gene, 57:267-272).

15.1.2. PLASMID AND YEAST STRAIN CONSTRUCTION

The coding region of human fchd540 was amplified by PCR and cloned in frame into pGBT9 (Bartel et al., 1993, Cellular Interactions in Development. pp. 153-159) resulting in plasmid pGBT9-fchd540. pGBT9-fchd540 was transformed into two-hybrid screening strain HF7c and one resulting transformant was designated TB35.

15.1.3. TWO-HYBRID SCREENING

Two-hybrid screening was carried out essentially as described (Bartel et al., 1993, *supra*) using TB35 as the recipient strain and a human breast two-hybrid library.

5

15.1.4 PAPER FILTER BETA-GALACTOSIDASE ASSAYS

The paper filter beta-galactosidase (beta-gal) assay was performed essentially as previously described (Brill et al., 1994, *Mol. Biol. Cell* 5: 297-312).

10

15.2 RESULTS

15.2.1 STRONG PHYSICAL INTERACTION OF RCHD534 AND FCHD540 MEASURED BY TWO-HYBRID ASSAY

15 The fchd540 coding sequence was amplified by PCR and cloned into pGBT9 creating a GAL4 DNA-binding domain-fchd540 fusion gene. The screening strain HF7c was transformed with this construct. The rchd534 coding sequence was cloned into pGAD424 (Bartel et al., 1993, *supra*) creating a GAL4 transcriptional activation domain-rchd534 fusion gene, which
20 was then used to transform strain Y187.

Yeast expression plasmids encoding the GAL4 DNA-binding domain either alone or fused in frame to fchd540, rchd534, *Drosophila* MAD, DPC4, or p53 were transformed into MAT α two-hybrid screening strain HF7c. Yeast expression
25 plasmids encoding the GAL4 transcriptional activation domain alone and GAL4 activation domain fusions to rchd534 and SV40 were transformed into MAT α two-hybrid screening strain Y187. p53 and SV40 interact with each other and should not interact with the experimental proteins. The HF7c transformants were
30 propagated as stripes on semisolid synthetic complete medium lacking L-tryptophan and the Y187 transformants were grown as stripes on semisolid synthetic complete medium lacking L-leucine. Both sets of stripes were replica plated in the form of a grid onto a single rich YPAD plate and the haploid
35 strains of opposite mating types were allowed to mate overnight at 30°C. The yeast strains on the mating plate were then replica plated to a synthetic complete plate

09034286 030498
lacking L-leucine and L-tryptophan to select for diploids and incubated at 30°C overnight. Diploid strains on the synthetic complete plate lacking L-leucine and L-tryptophan were replica plated to a synthetic complete plate lacking L-leucine, L-tryptophan, and L-histidine to assay HIS3 expression and a paper filter on a synthetic complete plate lacking L-leucine and L-tryptophan. The next day the paper filter was subjected to the paper filter beta-galactosidase assay to measure expression of the lacZ reporter gene. HIS3 expression was scored after 3 days of growth at 30°C. The results are shown in Table 3.

The rchd534 fish protein was found to interact strongly with the fchd540 bait protein and not to interact with the rchd534, MAD, DPC4, p53, and GAL4 DNA binding domain bait proteins. This result demonstrated that rchd534 and fchd540 strongly physically interact with each other with significant specificity.

15.2.2 IDENTIFICATION OF PROTEINS THAT PHYSICALLY INTERACT WITH FCHD540

The fchd540 coding sequence was amplified by PCR and cloned into pGBT9 (Bartel et al., 1993, *supra*) creating a GAL4 DNA-binding domain-fchd540 fusion gene. HF7c was transformed with this construct resulting in strain TB35. TB35 grew on synthetic complete medium lacking L-tryptophan but not on synthetic complete medium lacking L-tryptophan and L-histidine demonstrating that the GAL4 DNA-binding domain-fchd540 fusion does not have intrinsic transcriptional activation activity.

TB35 was transformed with the human breast two-hybrid library and 5 million transformants were obtained. The transformants were plated on synthetic complete medium lacking L-leucine, L-tryptophan, and L-histidine and yeast colonies that both grew on synthetic complete medium lacking L-leucine, L-tryptophan, and L-histidine and expressed the beta-galactosidase reporter gene were identified. The 30 strains with the strongest beta-galactosidase induction were characterized. Library plasmids were isolated from these

strains, and the 5' ends of all of the cDNA inserts were sequenced.

15.2.3 RETRANSFORMATION AND SPECIFICITY TESTING OF TCHV03A AND TCHVR4A

5

Two of the plasmids that encoded the strongest interactors were found to contain rchd534 cDNAs. Plasmid tchv03A was found to encode amino acids 17-235 of rchd534 and plasmid tchvR4A was found to encode amino acids 25-235 of rchd534.

10

It was confirmed that these rchd534 cDNAs encode proteins that physically interact specifically with fchd540. Yeast expression plasmids encoding the GAL4 DNA-binding domain either alone or fused in frame to fchd540, rchd534, Drosophila MAD, DPC4, and p53 were transformed into MAT α two-hybrid screening strain HF7c. Yeast expression plasmids encoding the GAL4 transcriptional activation domain (GAL4 AD) alone and GAL4 activation domain fusions to tchv03a, tchvR4A and SV40 were transformed into MAT α two-hybrid screening strain Y187. p53 and SV40 interact with each other and should not interact with the experimental proteins. The HF7c transformants were propagated as stripes on semi-solid synthetic complete medium lacking L-leucine. Both sets of stripes were replica plated in the form of a grid onto a single rich YPAD plate and the haploid strains of opposite mating types were allowed to mate overnight at 30°C. The yeast strains on the mating plate were then replica plated to a synthetic complete plate lacking L-leucine and L-tryptophan to select for diploids and incubated at 30°C overnight. Diploid strains on the synthetic complete plate lacking L-leucine and L-tryptophan were replica plated to a synthetic complete plate lacking L-leucine, L-tryptophan, and L-histidine to assay HIS3 expression and a paper filter on a synthetic complete plate lacking L-leucine and L-tryptophan. The next day the paper filter was subjected to the paper filter beta-galactosidase assay to measure expression of the lacZ reporter gene. HIS3 expression was scored after 3 days of growth at 30°C. The results are shown in the table below.

15

20

25

30

35

The strength or absence of physical interaction between each combination of test proteins is listed. Strong interactions are defined as interactions that cause the activation of both the HIS3 and lacZ reporter genes.

5

10

15

20

25

30

35

003428 030498
964060 98246060

TABLE 3

		cDNA-GAL4 Activation Domain Fusion Tested				
		rchd534	tchv03A	tchvR4A	SV40	GAL4 AD alone
5	GAL4 DNA-Binding Domain Fusions					
10	fchd540	Strong	Strong	Strong	None	None
	rchd534	None	None	None	None	None
	Dros. MAD	None	None	None	None	None
	DPC4	None	None	None	None	None
	p53	None	None	None	Strong	None
15	GAL4 DNA-Binding Domain alone	None	None	None	None	None

20 The tchv03A and tchvR4A fish proteins were found to interact strongly with the fchd540 bait protein and to not interact with the rchd534, MAD, DPC4, p53, and GAL4 DNA binding domain bait proteins. These results confirm the result that the rchd534 and fchd540 proteins interact strongly with each other.

25

15.3 FURTHER ANALYSIS OF RCHD534 AND FCHD540 FUNCTION

The significance of the rchd534/fchd540 protein interaction was confirmed by examination of their expression and activity in human cells and animal models.

30

15.3.1 CHROMOSOMAL LOCALIZATION

35 The rchd534 gene was localized to chromosome 15 and the fchd540 gene was localized to chromosome 18, regions of the human genome that contain other MAD homologues. These regions of the human genome have also been implicated in the pathogenesis of several human malignancies.

15.3.2 TISSUE EXPRESSION PATTERNS

The expression patterns were examined using in situ hybridization techniques. Fluorescently labeled DNA probes of both the rchd534 and fchd540 genes were used to probe human carotid endarterectomy samples. The expression of rchd534 and fchd540 was specific to endothelial cells lining the luminal surface of the carotid artery. In addition, a rabbit polyclonal antiserum generated against the rchd534 gene product prominently and selectively stained the endothelium present in large vessels such as human coronary arteries as well as smaller vessels present within human myocardium. Neither gene showed expression in any other cell type present in the arterial tissue sample, including smooth muscle cells and macrophages.

Expression patterns of both genes were also examined in response to certain stimulus. Both genes are selectively upregulated under the steady laminar shear stress (LSS) paradigm, but not under the turbulent shear stress paradigm or in response to stimulus by the cytokines rhIL-1 β , TNF α , IFN γ or active TGF β as measured in HUVEC cells. Thus, the rchd534 and the fchd540 genes appear to be selectively responsive to a LSS stimulus, manifesting no response to a non-laminar fluid mechanical stimulus, nor any other humoral stimuli tested. Thus, given that these two genes are: (1) localized to a region of the human genome that has been implicated in the pathogenesis of several human malignancies; (2) specifically expressed in a cell-type that is found only in vascular tissue, including atherosclerotic plaques; (3) up-regulated under the steady laminar shear stress cardiovascular disease paradigm; and (4) specifically inhibit TGF- β signalling indicate that rchd534 and fchd540 are excellent and specific targets for therapeutic intervention in the treatment of fibroproliferative and oncogenic disorders including tumor growth and vascularization.

35

15.3.3. CELLULAR LOCALIZATION

The cellular localization of the rchd534 and fchd540 proteins in bovine aortic endothelial cells (BAECs) was examined in relationship to other proteins involved in the TGF- β signalling pathway. In all experiments, the rchd534 and fchd540 proteins were located in the cytoplasm. MADR2 was located in the cytoplasm when transfected alone and in the nucleus when co-transfected with activated T β RI or when TGF- β was added to the culture medium. Co-transfection of rchd534 or fchd540 with MADR2 prevented the localization of MADR2 in the nucleus in response to TGF- β signalling.

The ability of the rchd534 and fchd540 proteins to interact with themselves and with other protein members of the TGF- β signalling pathway (MADR1, MADR2, DPC4, Tbr1, TSR1, ActR1b, ALK3, ALK6), was tested using this co-immunoprecipitation method. Each gene was transfected alone and in various combinations with other TGF- β pathway genes in either 293 cells or BAECs. The rchd534 protein formed homodimers in 293 cells and BAECs. The fchd540 protein did not form homodimers in 293 cells or BAECs. As mentioned above, the rchd534 and fchd540 proteins formed heterodimers in 293 cells and BAECs. This interaction is about 50 fold

stronger in BAECs than 293 cells based on equal amounts of protein. However, the rchd534-fchd540 protein interaction was significantly less avid than the rchd534 protein's interaction with itself.

- 5 The rchd534 protein interacted with MADR1, MADR2, and DPC4 in 293 cells and BAECs. The strength of MADR1 and MADR2 interactions was about the same between 293 cells and BAECs and much greater in BAECs for DPC4. The fchd540 protein interacted very weakly with MADR1, MADR2, and DPC4 in 293
- 10 cells. The rchd534 protein interacted strongly with activated forms of T β RI and ActRI and weakly with activated ALK6 in 293 cells. The fchd540 protein interacted strongly with activated T β RI and ALK6 receptors, and weakly with activated forms of TSRI, ALK3, and ActRIb in 293 cells.
- 15 Thus, in addition to the interaction of the rchd534 and fchd540 proteins, the interaction of the rchd534 protein with itself, as well as the interaction of the rchd534 protein and the fchd540 protein with the other proteins in the TGF- β pathway described above are excellent targets for therapeutic
- 20 intervention.

15.3.5 EFFECT OF EXPRESSION ON TGF-B SIGNALLING

- The effect of both rchd534 and fchd540 on the TGF- β signalling pathway was tested in vitro. Primary BAECs were
- 25 transfected with a construct called p3TP-Lux, containing a TGF- β responsive promoter fused to a reporter gene (Wrana et al., 1994, Nature 370: 341-347). The rchd534 gene or the fchd540 gene in pCI expression vectors (Promega) was transfected with and without MADR1 (pCMV5MADR1-Flag, Hoodless
- 30 et al. 1996 Cell 85: 489-500) or MADR2 (pCMV5MADR2-Flag, Eppert et al. 1996 Cell 86: 543-552). The TGF- β response was induced 20-fold by either MADR1 or MADR2. Co-expression of either rchd534 or fchd540 completely eliminated this induction. Thus, the rchd534 and fchd540 proteins inhibited
- 35 MADR1- and MADR2-mediated TGF- β signalling in endothelial cells. To confirm the specificity of this inhibitory effect, site specific mutants of both rchd534 or fchd540 were

constructed, based on known mutations identified in *Drosophila* homologues, that would be predicted to disrupt MAD-like signaling functions (Sekelsky et al., 1995, Genetics 139:1347-58; Raftery, 1995, Genetics 139:241-54; Newfeld et al., 1996, Development 122:2099-108; Wiersdorff et al., 1996, Development 122:2153-62). Unlike wild type *rchd534* and *fchd540*, these mutant proteins were unable to inhibit the activation of the p3TP promoter in response to TGF- β . The expression levels of the mutant and wild-type proteins were comparable indicating the loss of function was not due to secondary instability.

Interestingly, *Smad3*, the *C. elegans* homolog to *MAD3* which also functions in TGF β signalling is over 90% identical to *Smad2*, the *C. elegans* *MAD2* homolog, in the MH2 domain. Although this has not yet been directly investigated, it is likely that *Smad7*, the *C. elegans* homolog of the *fchd540* gene, may function similarly to its inhibition to prevent association and activation of *Smad3* by the TGF β receptor, that is, to inhibit the phosphorylation of *Smad3* and its association with protein components of the TGF- β signalling pathway.

These results further demonstrate that the interactions of either the *rchd534* protein or the *fchd540* protein with *MADR2* or with activated T β R1 are excellent targets for therapeutic intervention. As described above, the expression of *rchd534* and *fchd540* is specific, within arterial tissue, to endothelial cells. Accordingly, the *rchd534* and *fchd540* genes may be targets for intervention in a variety of inflammatory and fibroproliferative disorders that involve endothelial cells, including, but not limited to, cancer angiogenesis, inflammation, and fibrosis.

16. EXAMPLE: THE RCHD534-LONG PROTEIN

As described below, the rchd534 gene was discovered to encode two spliceoforms. The short spliceoform, and the rchd534 protein it encodes, are described in detail in Section 9, above. A second novel spliceoform, encoding a novel longer protein designated rchd534-long, is described in detail in the subsections below.

16.1 IDENTIFICATION AND CHARACTERIZATION OF
THE RCHD534-LONG SPLICEOFORM AND PROTEIN

10 A human heart cDNA library (Stratagene, LaJolla, CA) was screened with a probe containing nucleotides 400-700 of the fchd540 (see FIG.10) under the following hybridization conditions: hybridization overnight at 65°C, washing with 2xSSC and 0.1% SDS for 20 minutes at room temperature, 15 followed two washes with 0.2xSSC and 0.1%SDS for 20 minutes at 65°C. A positive clone was found to encode a novel protein, related to the rchd534 protein, that was designated the rchd534-long protein. The rchd534-long protein, like the rchd534 protein, has an MH2 domain. In addition, the 20 rchd534-long protein has an MH1 domain not present in rchd534. The original clone isolated from the Stratagene heart library contained incorrect sequence, including two stop codons, between the MH1 and MH2 domain coding regions. Therefore, cDNA prepared from human heart mRNA (Clontech, 25 Palo Alto, CA) was used as template for the PCR reaction to isolate the correct cDNA sequence in the region spanning the MH1 and MH2 domain coding regions. The following two pairs of nested primers were used:

Pair 1: A) 5'-~~GAGGCTGCGGCCGCTCEGAAGTCC~~-3'
B) 5'-CTCCGCCGGGGCCGCCACTATCT-3'

a Pair 2: A) 5'-CCGGGACGCAGTGGGACAG-3' (SEQ ID NO: 23)
a B) 5'-CGGGGAGTTGACGAAGATGG-3' (SEQ ID NO: 24)

35 The nucleotide sequence of the PCR amplification product was determined and confirmed to be accurate by sequence analysis of several amplification products from both

Clontech human lung cDNA, and heart mRNA that was obtained from Clontech and then reverse-transcribed into cDNA. The incorrect sequence in the original clone obtained from the Stratagene heart library was replaced by a PCR amplification product containing the correct sequence.

This correct cDNA encoding the rchd534-long protein was cloned into the TA cloning vector (Invitrogen) to create plasmid pHL6TA1A, which was deposited with the American Type Culture Collection on February 6, 1998 as Accession No.

209615. The cDNA sequence of the rchd534-long spliceoform encoding the entire rchd534-long protein is shown in FIG.9. The rchd534-long nucleotide sequence is 93% identical to the nucleotide sequence of the mouse SMAD6 gene (Imamura et al., 1997, Nature 389: 622-626). The rchd534-long sequence was reported in Hata et al., 1998, Genes and Development 12: 186-197.

The domains of the rchd534-long protein are shown in schematic form and compared with the domains of the rchd534 protein in FIG.11. The rchd534-long protein contains an MH1 domain, a spacer region, and an MH2 domain. The rchd534-long protein contains an N-terminal 273 amino acids (from Met-1 to Glu-273) which are not present in the rchd534 protein. This N-terminal region contains an MH1 domain that is more highly homologous to the MH1 domain of fchd540 than to the MH1 domain of other MAD proteins.

The amino acid sequence of rchd534-long protein from amino acid Ser-274 to the C-terminal amino acid Arg-496 are identical to amino acids Ser-13 to the C-terminal amino acid Arg-235 of the rchd534 protein. This region comprises an MH2 domain, from Pro-328 to Arg-496 in rchd534-long, and from Pro-67 to Arg-235 in rchd534. The rchd534 protein contains a 12 amino acid sequence at the N-terminus, from Met-1 to Lys-12, which is not present in the rchd534-long protein.

16.2. TGF- β SIGNALLING INHIBITORY
ACTIVITY OF THE RCHD534-LONG PROTEIN

The activity of the rchd534-long protein was tested using the TGF- β responsive reporter system described in Section 15.3.5, above for the rchd534 and fchd540 proteins.

HEPG2, 293 and BAEC cells were transfected with a construct called p3TP-Lux, containing a TGF- β responsive promoter fused to a reporter gene (Wrana et al., 1994, Nature 370: 341-347). The rchd534-long gene in pCI expression vectors (Promega) was transfected with and without MADR2. Luciferase activity produced from the reporter construct was measured relative to an internal control (secreted alkaline phosphatase expression). The results obtained in 293 (human embryonic kidney) cells are summarized in Table 4 below. A "+" indicates that the specified construct, plasmid, or gene was present in the cell line assayed; whereas a "-" indicates that it was absent.

TABLE 4

	3TP-LUX	TGF-bRI	pCI	MADR2	fchd540	rchd534	rchd534-long	Relative Luciferase Activity
20	+	-	-	-	-	-	-	10
	+	+	-	-	-	-	-	48
	+	+	+	-	-	-	-	40
	+	+	-	+	-	-	-	107
25	+	+	-	-	+	-	-	2
	+	+	-	-	-	+	-	10
	+	+	-	-	-	-	+	2
	+	+	+	+	-	-	-	104
30	+	+	-	+	-	-	-	124
	+	+	-	+	+	-	-	13
	+	+	-	+	-	+	-	59
	+	+	-	+	-	-	+	18

The TGF- β response, both un-induced by MADR2 and induced by MADR2 overexpression, was inhibited by expression of rchd534-long, as well as by rchd534 and fchd540. Similar

inhibition of TGF- β signalling by the rchd534-long protein was observed in each of the several different cell lines.

17. EXAMPLE: ANTISENSE AND RIBOZYME MOLECULES FOR INHIBITION OF RCHD534 AND FCHD540 EXPRESSION

The principles presented in Section 5.6.1.1, above, can be used to design oligonucleotides for use in inhibiting the expression of target genes, such as the rchd534 or fchd540 genes.

10

17.1. ANTISENSE MOLECULES

The following antisense molecules can be used to inhibit translation of the rchd534 protein:

- a₁₅ a) 5'-CATTTTCATTTTCATACAA-3' ^(SEQ ID NO:25) which is complementary to nucleotides -14 to +3 of in FIG.8.
- a b) 5'-CATTTTCATTTTCATACAATATATG-3' ^(SEQ ID NO:26) which is complementary to nucleotides -20 to +3 in FIG.8.
- a₂₀ c) 5'-CATTTTCATTTTCATACAATATATGGCCTTT-3' ^(SEQ ID NO:27) which is complementary to nucleotides -26 to +3 in FIG.8.
- a₂₅ d) 5'-CATTTTCATTTTCATACAATATATGGCCTTTTGTGGC-3' ^(SEQ ID NO:28) which is complementary to nucleotides -32 to +3 in FIG.8.
- a e) 5'-GGACATTTTCATTTTCATACAATATATGGCCTTTTGT-3' ^(SEQ ID NO:29) which is complementary to nucleotides -29 to +6 in FIG.8.
- a₃₀ f) 5'-TTCATTTTCATACAATATATGGCCTTTTGT-3' ^(SEQ ID NO:30) which is complementary to nucleotides -29 to -1 in FIG.8.
- a g) 5'-TCATACAATATATGGCCTTTTGT-3' ^(SEQ ID NO:31) which is complementary to nucleotides -29 to -7 in FIG.8.
- a₃₅ h) 5'-AATATATGGCCTTTTGT-3' ^(SEQ ID NO:32) which is complementary to nucleotides -29 to -13 in FIG.8.

The following antisense molecules can be used to inhibit translation of the rchd534-long protein:

- Q a) 5'-CATACGATATCCTTTGGCGCCAGGGG-3' ^(SEQ ID NO:33) which is
5 complementary to nucleotides -23 TO +3 in FIG.9.
- a b) 5'-GGACCTGAACATACGATATCCTTTGGCGCCAGGGG-3' ^(SEQ ID NO:34) which is
complementary to nucleotides -23 TO +12 in FIG.9.
- Q 10 c) 5'-CATACGATATCCTTTGGCGCCAGGGGTGGGGGGG-3' ^(SEQ ID NO:35) which is
complementary to nucleotides -31 TO +3 in FIG.9.

The following antisense molecules can be used to inhibit translation of the fchd540 protein:

- a a) 5'-CATGCGGGGCGAGGAGG-3' ^(SEQ ID NO:36) which is complementary to
nucleotides -14 to +3 of fchd540 in FIG.10.
- a b) 5'-CATGCGGGGCGAGGAGGCGAGGA-3' ^(SEQ ID NO:37) which is complementary
20 to nucleotides -20 to +3 of fchd540 in FIG.10.
- a c) 5'-CATGCGGGGCGAGGAGGCGAGGAGAAAAG-3' ^(SEQ ID NO:38) which is
complementary to nucleotides -26 to +3 of fchd540 in
FIG.10.
- 25 a d) 5'-CATGCGGGGCGAGGAGGCGAGGAGAAAAGTCGTTT-3' ^(SEQ ID NO:39) which is
complementary to nucleotides -32 to +3 of fchd540 in
FIG.10.
- Q 30 e) 5'-GAACATGCGGGGCGAGGAGGCGAGGAGAAAAGTCG-3' ^(SEQ ID NO:40) which is
complementary to nucleotides -29 to +6 of fchd540 in
FIG.10.
- a f) 5'-GCGGGGCGAGGAGGCGAGGAGAAAAGTCG-3' ^(SEQ ID NO:41) which is
35 complementary to nucleotides -29 to -1 of fchd540 in
FIG.10.

g) 5'-CGAGGAGGCGAGGAGAAAAGTCG-3' ^(SEQ ID NO: 42) which is complementary to nucleotides -29 to -7 of fchd540 in FIG.10.

h) 5'-GGCGAGGAGAAAAGTCG-3' ^(SEQ ID NO: 43) which is complementary to nucleotides -29 to -13 of fchd540 in FIG.10.

17.2. RIBOZYME MOLECULES

The central, catalytic portion of a hammerhead ribozyme molecule consist of the following sequence:

5'-CAAAGCNGNXXXXNCNGAGNAGUC-3' ^(SEQ ID NO: 44)

wherein the 5'-proximal CA bases hybridize to a complementary 5'-UG-3' in the target mRNA. The first four underlined bases form a stem by base pairing with the second set of underlined bases, with the intervening bases, shown as X's, forming a non-pairing loop. In order to hybridize to a target mRNA, a hammerhead ribozyme contains additional bases flanking each end of the central segment shown above. The 5' ribozyme flanking segment is complementary to the respective flanking sequences immediately 3' to the target UG; and the 3' flanking segment is complementary to the respective flanking sequence beginning two bases upstream of the target U, and extending 5'-ward (in effect, skipping the first base upstream of the target U). Cleavage occurs between first and second bases upstream of (i.e., 5' to) the U in the target 5'-UG-3' site.

The following ribozyme molecules can be used to inhibit translation of the rchd534 protein:

a) 5'-GGUGGAGCCCCAGGGCAUUAACCUCAAAGCNGNXXXXNCNGAGNAGUCGUGG
GCAAGGUGGGCACUCAGGUGGG-3' ^(SEQ ID NO: 45) which will cleave the short spliceoform rchd534 mRNA between nucleotides 716 and 717 in FIG.8.

b) 5'-GUGUCUCUAUGGGUUUGCCCAAAGCNGNXXXXNCNGAGNAGUCUCUGGACA
UUUCAUUUCAUAC-3' ^(SEQ ID NO: 46) which will cleave the short

spliceoform rchd534 mRNA between nucleotides 1040 and 1041 in FIG.8.

The following ribozyme molecules can be used to inhibit translation of the rchd534 protein or the rchd534-long protein:

- a) 5'-GGCCCUCUCGCCGUCGGGCUCCUUGCUGAGCAAAGCNGNXXXXXNCNGAGNA
GUCGAUGCCGAAGCCGAUCUUGCUGCGCG-3' ^(SEQ ID NO: 47) which will cleave the
short spliceoform rchd534 mRNA between nucleotides
1421 and 1422 in FIG.8; and the rchd534-long
spliceoform mRNA and between nucleotides 1327 and 1328
in FIG.9.

The following ribozyme molecules can be used to inhibit translation of the rchd534-long protein:

- a) 5'-CGACUUCGCCAAAGUCGCCGCAAAGCNGNXXXXXNCNGAGNAGUCCAGCCCC
GAGCGTTTGGACCTG-3' ^(SEQ ID NO: 48) which will cleave between
nucleotides +178 and +179 in FIG. 9.

The following ribozyme molecules can be used to inhibit translation of the fchd540 protein:

- a) 5'-CGUUUGCCUGCUAAGGAGCGAACAAGCNGNXXXXXNCNGAGNAGUCGAUGU
UUCUUUGUGAGUCGGGCGCCG-3' ^(SEQ ID NO: 49) which will cleave the
fchd540 mRNA between nucleotides -53 and -52 in
FIG.10.
- b) 5'-CGCCGGACGAGCGCAGAUCGUUUGGUCCUGAACAAAGCNGNXXXXXNCNGAG
NAGUCCGGGGCGAGGAGGCGAGGAGAAAAGUCG-3' ^(SEQ ID NO: 50) which will
cleave the fchd540 mRNA between nucleotides -1 and +1
in FIG.10.
- c) 5'-GGAGUAAGGAGGGGGGGGAGACUCUAGUUCGCAAAGCNGNXXXXXNCNGAGN
AGUCAGUCGGCUAAGGUGAUGGGGGUUGCAGCACACC-3' ^(SEQ ID NO: 51) which will
cleave the fchd540 mRNA between nucleotides +602 and
+603 in FIG.10.

[illegible]

10

RCHD528 B-21382

20

FCHD534 B-21459

30

FCHD502SJ 69982

PEDC-114894.1

The following microorganisms were deposited with the American Type Culture Collection (ATCC), Rockville, Maryland, on February 9, 1996, and assigned the indicated accession numbers:

5	<u>Microorganism</u>	<u>ATCC Accession No.</u>
	FCHD528A	69985
	FCHD528B	69986
	FCHD528C	69987

10 The following microorganism was deposited with the American Type Culture Collection (ATCC), Rockville, Maryland, on February 7, 1996, and assigned the indicated accession number:

	<u>Microorganism</u>	<u>ATCC Accession No.</u>
15	pFCHD540	69984

The following plasmid, encoding the rchd534-long protein, was deposited with the American Type Culture Collection (ATCC), Rockville, Maryland, on February 6, 1998, and assigned the indicated accession number:

<u>Microorganism</u>	<u>ATCC Accession No.</u>
pHL6TA1A	209615

25 The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various
30 modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

35